arXiv:1202.5754v2 [math.GT] 18 Apr 2012

Preprint (2012)

HIGHER ORDER GENERALIZATION OF FUKAYA'S MORSE HOMOTOPY INVARIANT OF 3-MANIFOLDS I. INVARIANTS OF HOMOLOGY 3-SPHERES

TADAYUKI WATANABE

Abstract. We give a higher order generalization of Fukaya's Morse homotopy theoretic approach to 2-loop Chern–Simons perturbation theory. In this paper, we construct a sequence of invariants of homology 3-spheres with values in a space of trivalent graphs (Jacobi diagrams) by using Morse homotopy theory.

1. **Introduction**

K. Fukaya constructed in [Fuk96] a 3-manifold invariant by using his Morse homotopy theory, which is defined by considering several Morse functions simultaneously. Fukaya's invariant is conjectured to coincide with the 2-loop part of Chern–Simons perturbation theory of Axelrod–Singer and Kontsevich ([AS, Ko]). The universal form of the 3-manifold invariant of Chern–Simons perturbation theory has values in a space of graphs and is known to be universal among Ohtsuki's finite type invariants if restricted to homology spheres ([Oh96, KT, Les]). Fukaya's invariant is defined by counting graphs in a manifold whereas Chern–Simons perturbation theory invariant is defined by an integration over the configuration space of a 3-manifold.

Behind the work of [Fuk96], there is a work of physics due to Witten that the open string theory on the cotangent bundle of a 3-manifold is equivalent to the Chern–Simons perturbation theory ([Wi]). Indeed, in [FO], it has been proved mathematically that the 0-loop open string theory on the cotangent bundle of a manifold is equivalent to the Morse homotopy theory, which had been independently developed by Betz–Cohen ([BC]) and by Fukaya ([Fuk93]). There is also a discussion in [Fuk93, Fuk96] about a relationship between the higher genus open string theory and Chern–Simons perturbation theory.

In [Fuk96], Fukaya considered triads of Morse functions (f_1, f_2, f_3) on M and for pairs (M, ζ_i), $i = 1, 2$, of a closed 3-manifold M and acyclic flat $SU(2)$-bundles ζ_i, he defined some number $\hat{Z}_2(M, f_1, f_2, f_3; \zeta_i)$. He showed that the difference $\hat{Z}_2(M, f_1, f_2, f_3; \zeta_1) - \hat{Z}_2(M, f_1, f_2, f_3; \zeta_2)$ depends only on the equivalence class of (M, ζ_1, ζ_2). The essential points in his construction may be the following.

Date: April 19, 2012.

2000 *Mathematics Subject Classification.* 57M27, 57R57, 58D29, 58E05.

(1) Several Morse functions are considered at once and graph-shaped flows along the gradients are counted. The moduli spaces of the flows form compact 0-dimensional manifolds for a generic triad of Morse functions.
(2) Graphs with an edge separated in the middle by critical points (as in Figure 1) are considered and a linear combination of the numbers of (possibly separated) graph-flows is considered. Here the coefficients are endomorphisms between tensor products of the Morse complexes (with twisted coefficients).
(3) For each coefficient in the linear combination, combinatorial propagators (or chain contractions) of Morse complexes are composed and then the trace is taken to obtain a number.

The aim of this paper is to construct graph-valued invariants of homology 3-spheres, i.e., closed 3-manifolds M with $H_*(M) \cong H_*(S^3)$, by using Morse homotopy theory, as a higher order generalization of [Fuk96]. In this paper, we generalize the idea of Fukaya to graphs with loop numbers (the first Betti numbers) ≥ 2 for homology 3-sphere M with the trivial connection. To give an explicit formula for our invariant for all orders, we introduce an appropriate graph complex for Morse homotopy theory being based on Kontsevich's graph complex in [Ko] and formulate the formal cancellation mechanism among different graphs, that will be necessary in the proof of invariance of our invariant, in the graph complex. The principal term $Z_{2k,3k}$ of our invariant is defined by substituting the numbers of graph-flows into cycles in our graph complex.

The proof that our invariant is well-defined is done by a cobordism argument for a 1-parameter family of smooth functions on M without higher singularities. The main point in the cobordism argument is that the relations among the numbers of moduli spaces of graph-flows in 1-parameter family are similar to the relations in the graph complex. In this paper, we consider graphs for all orders, so we attempt to give full details of the structure of a manifold with corners of a moduli space and of arguments of orientations etc.

The moduli space of graph-flows will be described as the transversal intersections of several submanifolds of the configuration space of M or the direct product of the configuration space of M with $[0, 1]$. To make such definition easier, we will not use, in the cobordism argument, the argument of time dependent trajectories of [Fuk96]. Instead, we confirm the invariance one at a time by using a similar bifurcation analysis as in [Ce].

Also, in this paper, we consider only the trivial connection contribution and define an invariant not by taking the difference as in [Fuk96]. To obtain an invariant not by taking the difference, we need to introduce an anomaly correction term appropriately. In [Fuk96], the existence of such a correction term in the 2-loop case was a conjecture. In this paper, we introduce an anomaly term $Z_{2k,3k}^{\mathrm{anomaly}}$ by taking some linear combination of the numbers of infinitesimal graph-flows in a configuration space bundle over a compact 4-manifold having M as a boundary. The key point for the correction term to work well is the spin cobordism invariance of the anomaly term $Z_{2k,3k}^{\mathrm{anomaly}}$. Namely, the spin cobordism invariance allows us to define an analogue of the signature defect, which includes $Z_{2k,3k}^{\mathrm{anomaly}}$ instead of the relative L-class in the sigunature defect, and it gives the desired correction term.

There are analogues of our graph complex and the moduli spaces of graph-flows for circle-valued Morse theory. The generalization to circle-valued Morse function would give an invariant of 3-manifolds with the first Betti number 1. We plan to discuss this in [Wa2].

Finally, we mention M. Futaki's work [Fut]. Futaki discovered in [Fut] some singular phenomena that are missed in [Fuk96]. In [Fuk96], the coefficients in the linear combination of graphs are defined by contracting holonomies along graph-flows by $\mathfrak{su}(2)$-invariant tensors. However, Futaki observed with a concrete calculation that when the dumbbell graph contribution is nontrivial (homology 3-sphere with the trivial connection is not the case), the holonomy matrix will suddenly jump at the point on which a trivalent vertex passes through a critical point and thus the invariance fails. Since we construct an invariant via an intersection theory considering only the trivial connection contribution, the coefficients in the linear combination in our definition can be given without using holonomy matrix, so the same problem do not occur. (See also Remark 3.1.)

Organization. The organization of the present paper is as follows. In chapter 2 we introduce our graph complex and construct a cycle in it. The cycle condition in the graph complex is a formal description of the cancellation of singular flows that appear in the proof of invariance of our invariant. In chapter 3, we give definitions of Fukaya's moduli spaces of graph-flows and the moduli space of infinitesimal graph-flows, and then consider their compactifications and orientations. Also, the definition of our invariant will be given and the main theorem, which says that our invariant is actually an invariant, is stated (Theorem 3.32). In chapter 4, we consider the compactification and orientation for the moduli space of graph-flows in 1-parameter family of smooth functions. In chapter 5, in accordance with the contents in previous chapters, we prove the invariance of our invariant by a cobordism argument. For each of the four types of bifurcations that may occur generically, we confirm the invariance one at a time.

Convention. We denote by $C^r(M)$ the space of C^r functions $f : M \to \mathbb{R}$ on a compact manifold M for sufficiently large r and we equip $C^r(M)$ the C^r-topology. For a C^r function f on a manifold M, we denote by $\Sigma(f)$ the subset of M of critical points of f. Let $\Sigma^1(f)$ (resp. $\Sigma^2(f)$) denote the subset of $\Sigma(f)$ consisting of Morse singularities (resp. birth/death points, or A_2-singularities). For a Morse function f, a critical point p of f and a metric μ on a manifold, we denote by $\mathscr{D}_p(f) = \mathscr{D}_p(f;\mu)$ (resp. $\mathscr{A}_p(f) = \mathscr{A}_p(f;\mu)$) the descending manifold (resp. ascending manifold) of (f,μ) at p. We denote by $\Gamma(E)$ the space of sections of a fiber bundle $E \to B$. For a sequence of submanifolds $A_1, A_2, \ldots, A_r \subset W$ of a smooth Riemannian manifold W, we say that the intersection $A_1 \cap A_2 \cap \cdots \cap A_r$ is *transversal* if for each point x in the intersection, the subspace $N_xA_1 + N_xA_2 + \cdots + N_xA_r \subset T_xW$ spans the direct sum $N_xA_1 \oplus N_xA_2 \oplus \cdots \oplus N_xA_r$. The convention for orientations of manifolds is given in Appendix A.



Figure 1.

2. Graph complex

2.1. **Graphs.** By a *graph*, we mean a finite graph with each edge oriented, i.e., an ordering of the boundary vertices of an edge is fixed, and without self-loop. For a graph Γ, let $\mathrm{In}(\Gamma)$ (resp. $\mathrm{Out}(\Gamma)$) be the set of univalent vertices of Γ that are inputs (resp. outputs). Let $W(\Gamma)$ be the set of vertices of valence ≤ 2 ("white vertex"). Let $B(\Gamma)$ be the set of vertices of valence ≥ 3 ("black vertex"). An *admissible graph* is a graph such that

1. $|\mathrm{In}(\Gamma)| = |\mathrm{Out}(\Gamma)|$,
2. a bijection $\rho : \mathrm{In}(\Gamma) \to \mathrm{Out}(\Gamma)$ is fixed, and
3. each bivalent vertex has one incoming and one outgoing edges.

A *compact edge* of Γ is an edge connecting two black vertices. A *separated edge* of Γ is a pair of edges $\{(a,x),(y,b)\}$ with $x,y \in B(\Gamma)$, $a \in \mathrm{In}(\Gamma)$, $b \in \mathrm{Out}(\Gamma)$ such that $b = \rho(a)$. A *broken edge* of Γ is a pair of edges $\{(x,a),(a,y)\}$ with $x,y \in B(\Gamma)$, $a \in W(\Gamma)$. A *broken separated edge* of Γ is either a triple $\{(a,b),(b,x),(y,c)\}$ or a triple $\{(a,x),(y,b),(b,c)\}$, with $x,y \in B(\Gamma)$, $a,b,c \in W(\Gamma)$ such that $c = \rho(a)$. Let $\mathrm{Comp}(\Gamma)$, $\mathrm{Se}(\Gamma)$, $\mathrm{Br}(\Gamma)$, $\mathrm{Se}'(\Gamma)$ be the set of compact, separated, broken, broken separated edges respectively. A *regular edge* of Γ is either a compact edge or a separated edge of Γ. Let $\mathrm{Reg}(\Gamma)$ be the set of regular edges of Γ. Let $E(\Gamma) = \mathrm{Comp}(\Gamma) \cup \mathrm{Se}(\Gamma) \cup \mathrm{Br}(\Gamma) \cup \mathrm{Se}'(\Gamma)$. See Figure 1.

A *labeled graph* is an admissible graph Γ equipped with bijections $\alpha : \{1,2,\ldots,n\} \to B(\Gamma)$ and $\beta : \{1,2,\ldots,\ell\} \to E(\Gamma)$. Let $C_*^{(i)} = (C_*^{(i)}, \partial^{(i)})$, $C_*^{(i)} = \mathbb{Z}^{P_*^{(i)}}$, $i = 1,2,\ldots,\ell$, be a sequence of finitely based acyclic chain complexes over $\mathbb{Z}$. For $\vec{C} = (C_*^{(1)},\ldots,C_*^{(\ell)})$, we define a *$\vec{C}$-colored graph* as a labeled graph $\Gamma = (\Gamma,\alpha,\beta)$ such that on each white vertex of $\beta(i)$ a basis vector $p \in C_*^{(i)}$ is attached for each i.

For each edge $e = \beta(i)$ in a $\vec{C}$-colored graph, we define its *degree* by

$$\deg(e) = \begin{cases} 1 & \text{if } e \in \mathrm{Comp}(\Gamma) \\ d(p) - d(q) & \text{if } e \in \mathrm{Se}(\Gamma), \\ & \quad p \in C_*^{(i)} \text{ on input, } q \in C_*^{(i)} \text{ on output} \\ 0 & \text{if } e \in \mathrm{Br}(\Gamma) \\ d(p) - d(q) - 1 & \text{if } e \in \mathrm{Se}'(\Gamma), \\ & \quad p \in C_*^{(i)} \text{ on input, } q \in C_*^{(i)} \text{ on output} \end{cases}$$

where $d(x)$ denotes the degree of x. We define the *degree* of a $\vec{C}$-colored graph by

$$\deg(\Gamma) = (\deg(\beta(1)), \dots, \deg(\beta(\ell))).$$

We will call a $\vec{C}$-colored graph with degree $\vec{\eta} = (\eta_1, \dots, \eta_\ell)$, with n vertices and with $|\mathrm{Reg}(\Gamma)| = m$ a *$\vec{C}$-colored graph of type* $(n, m, \vec{\eta})$. We define the *closure* $\widehat{\Gamma}$ of Γ as the graph obtained from Γ by identifying white vertices of each pair $(a, \rho(a))$. An example of a $\vec{C}$-colored graph of type $(4, 6, (1, 1, 1, 1, 1, 1))$ is given in Example 3.2.

2.2. **Graph complex.** Let $\mathscr{G}^0_{n,m,\vec{\eta}}(\vec{C})$ be the set of pairs (Γ, or), where

(1) Γ is a $\vec{C}$-colored graph of type $(n, m, \vec{\eta})$ with connected closure $\widehat{\Gamma}$,
(2) or is an orientation of the real vector space

$$\mathbb{R}^{B(\Gamma)} \oplus \bigoplus_{e \in \mathrm{Comp}(\Gamma)} \mathbb{R}^{H(e)},$$

where $H(e) = \{e_+, e_-\}$ is the set of "half-edges",

Let $\mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ be the vector space over $\mathbb{Q}$ spanned by $\mathscr{G}^0_{n,m,\vec{\eta}}(\vec{C})$, quotiented by the relation

$$(\Gamma, -\mathrm{or}) = -(\Gamma, \mathrm{or}).$$

The bijection α and the edge orientation of a graph Γ defines a canonical graph orientation or_Γ. In the following, we assume that all graphs are oriented by canonical orientation, namely, if a graph is oriented by the opposite orientation to the canonical one, then we represent the oriented graph by minus of the canonically oriented one.

Let $\mathscr{G}^{\mathrm{comp}}_{n,m,(1,\dots,1)}(\vec{C})$ be the subspace of $\mathscr{G}_{n,m,(1,\dots,1)}(\vec{C})$ spanned by graphs with only compact edges. The sequence of complexes $\vec{C}$ is unnecessary to represent a graph in $\mathscr{G}^{\mathrm{comp}}_{n,m,(1,\dots,1)}(\vec{C})$, so there are canonical isomorphisms between $\mathscr{G}^{\mathrm{comp}}_{n,m,(1,\dots,1)}(\vec{C})$ for different sequences $\vec{C}$. Identifying $\mathscr{G}^{\mathrm{comp}}_{n,m,(1,\dots,1)}(\vec{C})$ for all $\vec{C}$, we simply write

$$\mathscr{G}_{n,m} = \mathscr{G}^{\mathrm{comp}}_{n,m,(1,\dots,1)}(\vec{C}).$$

We define a linear map

$$d : \mathscr{G}_{n,m,\vec{\eta}}(\vec{C}) \to \bigoplus_{j=1}^{\ell} \mathscr{G}_{n-1,m-1,(\eta_1,\dots,\widehat{\eta_j},\dots,\eta_\ell)}(C_*^{(1)}, \dots, \widehat{C}_*^{(j)}, \dots, C_*^{(\ell)})$$

by

$$d(\Gamma, \mathrm{or}) = \sum_{e \in \mathrm{Comp}(\Gamma)} (\Gamma/e, \text{induced ori}),$$

where for $e_j = (u, v)$ (u, v are vertices) the induced orientation of Γ/e_j is formally given by $i\left(\frac{\partial}{\partial v}\right)(dv_1 \wedge \cdots \wedge dv_n) \wedge (de_1^+ \wedge de_1^-) \wedge \overset{j}{\check{\cdots}} \wedge (de_m^+ \wedge de_m^-)$. It follows from the identity $i(X)i(Y) + i(Y)i(X) = 0$ that $d \circ d = 0$. Also, we define another linear map

$$d' : \mathscr{G}_{n,m,\vec{\eta}}(\vec{C}) \to \bigoplus_{j=1}^{\ell} \mathscr{G}_{n,m-1,(\eta_1,\dots,\eta_j - 1,\dots,\eta_\ell)}(\vec{C})$$

by $\sum_{e\in \mathrm{Reg}(\Gamma)} d'_e$, where

d'_e 

$= -\sum_{r_i\in P_*^{(i)}}$ 

$(\beta(i)=e)$

d'_e 

$= \sum_{\substack{r_i\in P_*^{(i)}\\ d(r_i)=d(p_i)-1}}$ 

$+ \sum_{\substack{s_i\in P_*^{(i)}\\ d(s_i)=d(q_i)+1}} (-1)^{d(p_i)-d(s_i)}$ 

$(\beta(i)=e)$,

where $\mathrm{or}(d'_e\Gamma)$ is the naturally induced one. (Note that $d'\circ d'\neq 0$.) The reason for the signs in the formula will be explained in §3.8.

2.3. **The space $\mathscr{H}_n$.** We shall introduce some relations in the space of graphs to bring them closer to the moduli space of graph flows. Let $\partial^{(i)}(x,y)$, $x,y\in C_*^{(i)}$, denote the graph 

. Let Γ be a $\vec{C}$-labeled graph with $\beta(i)\in \mathrm{Se}(\Gamma)$ and with basis vectors x and $y\in C_*^{(i)}$ attached on the input and output white vertex respectively. To specify that Γ has such an edge, we will write $\Gamma=\Gamma(x,y)_i$. This notation enables us to express the graph obtained from Γ by replacing x and y with x' and y' respectively, as $\Gamma(x',y')_i$. We will write $\Gamma(\emptyset,\emptyset)_i$ the graph obtained from Γ by replacing the separated edge $\beta(i)$ with a compact edge.

Let

$$\mathscr{G}_{n,*,\vec{\eta}}(\vec{C})=\bigoplus_{m=0}^{\ell}\mathscr{G}_{n,m,\vec{\eta}}(\vec{C}),$$

$$\mathscr{H}_n(\vec{C})=\frac{\bigoplus_{\vec{\eta}\leq(2,\ldots,2)}\mathscr{G}_{n,*,\vec{\eta}}(\vec{C})\oplus\mathscr{G}_{n-1,*,\vec{\eta}}(\vec{C})}{(d+d')\mathscr{G}_{n,*,\Sigma(1,\ldots,1)}(\vec{C}),\partial\text{-relation},C\text{-relation}},$$

where

1. $\Sigma(1,1,\ldots,1)=(2,1,\ldots,1)+(1,2,\ldots,1)+\cdots+(1,1,\ldots,2)$, i.e., $\mathscr{G}_{n,*,\Sigma(1,\ldots,1)}=\mathscr{G}_{n,*,(2,1,\ldots,1)}\oplus\cdots\oplus\mathscr{G}_{n,*,(1,1,\ldots,2)}$,
2. ∂-relation is given by
$$\partial^{(i)}(p_i,r_i)*\Gamma(r_i,q_i)_i=\partial^{(i)}_{p_ir_i}\cdot\Gamma(r_i,q_i)_i$$
$$\Gamma(p_i,r_i)_i*\partial^{(i)}(r_i,q_i)=\partial^{(i)}_{r_iq_i}\cdot\Gamma(p_i,r_i)_i$$
$(\partial^{(i)}(p_i,r_i)*\Gamma(r_i,q_i)_i$ etc. is a composition of two graphs at the univalent vertices, on which r_i is attached. $\partial^{(i)}_{p_ir_i}$ etc. is the incidence coefficient of r_i in $\partial^{(i)}p_i$.),
3. C-relation is given by



$=$ 

.

We denote by $\langle\Gamma\rangle$ the equivalence class in $\mathscr{H}_n(\vec{C})$ represented by Γ. The relation $(d+d')\mathscr{G}_{n,*,\Sigma(1,\ldots,1)}(\vec{C})$ will correspond to the fact that the signed number of boundaries of a moduli space that is an oriented 1-manifold is 0.

2.4. **Cycles in $\mathscr{G}_{n,m}$.** In the following, we assume that $\ell = m$, i.e., $\vec{C} = (C_*^{(1)}, \dots, C_*^{(m)})$. In such a case, we write $\mathscr{G}_{n,m}(\vec{C})$ the subspace of $\mathscr{G}_{n,m,(1,\dots,1)}(\vec{C})$ spanned by graphs with no bivalent vertices.

We consider a linear combination

$$\gamma = \sum_{\Gamma \in \mathscr{G}_{n,m}(\vec{C})} a(\Gamma)\Gamma \in \mathscr{G}_{n,m}(\vec{C}), \quad a(\Gamma) \in \mathbb{Q},$$

where the sum is taken over all $\vec{C}$-colored graphs in $\mathscr{G}^0_{n,m}(\vec{C})$ of type $(n, m, (1, \dots, 1))$, each equipped with canonical orientation. So we consider all possible labellings and edge orientations. The d-cycle condition

$$d\gamma = 0 \Leftrightarrow \sum_{\Gamma \in \mathscr{G}_{n,m}(\vec{C})} a(\Gamma) d\Gamma = 0$$

is equivalent to a system of linear equations among coefficients $a(\Gamma)$. We call the linear system the $(*)$-*relation.* By considering $a(\Gamma)$ as formal variables, we get a universal cycle. Namely, we define the space $\mathscr{A}_{n,m}(\vec{C})$ to be the quotient space of $\mathscr{G}_{n,m}(\vec{C})$ by the $(*)$-relation and the *label change relation*, which identifies two labeled oriented graphs with isomorphic underlying graph when one is obtained from another by even swappings of labels or of edge orientations. We denote by $[\Gamma]$ the equivalence class in $\mathscr{A}_{n,m}(\vec{C})$ represented by Γ. Then we put

$$\widetilde{\gamma}_{n,m} = \sum_{\Gamma \in \mathscr{G}_{n,m}(\vec{C})} [\Gamma] \otimes \Gamma \in \mathscr{A}_{n,m}(\vec{C}) \otimes \mathscr{G}_{n,m}(\vec{C}).$$

Clearly, $\widetilde{\gamma}_{n,m}$ is a $(1 \otimes d)$-cycle.

We will need a $(d + d')$-cycle later[*]. To make $\widetilde{\gamma}_{n,m}$ a $(d+d')$-cycle, we define the ξ-*relation* in $\mathscr{A}_{n,m}(\vec{C})$ by

$$\sum_{\substack{x_i \in C_*^{(i)} \\ d(x_i) = d(q_i)+1}} \partial^{(i)}_{x_i p_i} \cdot [\Gamma(x_i, q_i)_i] + \sum_{\substack{y_i \in C_*^{(i)} \\ d(y_i) = d(p_i)-1}} \partial^{(i)}_{q_i y_i} \cdot [\Gamma(p_i, y_i)_i] - \delta_{p_i q_i} \cdot [\Gamma(\emptyset, \emptyset)_i] = 0,$$

for each $\Gamma(p_i, q_i)_i \in \mathscr{G}_{n,m,(\eta_1,\dots,\eta_m)}(\vec{C})$, $\eta_j = 1$ $(j \neq i)$, $\eta_i = 0$.

Lemma 2.1. *$\langle (1 \otimes (d + d'))\widetilde{\gamma}_{n,m} \rangle = 0$ in $\mathscr{A}_{n,m}(\vec{C})/(\xi$-relation$) \otimes \mathscr{H}_n(\vec{C})$.*

Proof. We have seen that $(1 \otimes d)\widetilde{\gamma}_{n,m} = 0$. So it suffices to show that $\langle (1 \otimes d')\widetilde{\gamma}_{n,m} \rangle = 0$. We show that $(1 \otimes d')\widetilde{\gamma}_{n,m}$ vanishes in $\mathscr{A}_{n,m}(\vec{C})/(\xi\text{-relation}) \otimes \mathscr{I}_n(\vec{C})$, where

$$\mathscr{I}_n(\vec{C}) = \bigoplus_{\vec{\eta} \leq (2,\dots,2)} \mathscr{G}_{n,*,\vec{\eta}}(\vec{C}) \,\big/\, \partial\text{-relation}, C\text{-relation}.$$

We denote by $\langle \Gamma \rangle_1$ the equivalence class of Γ in $\mathscr{I}_n(\vec{C})$. Since the left hand side of the identity of Lemma 2.1 lies in the image from $\mathscr{A}_{n,m}(\vec{C})/(\xi\text{-relation}) \otimes \mathscr{I}_n(\vec{C})$, it suffices to check the assertion in the space $\mathscr{A}_{n,m}(\vec{C})/(\xi\text{-relation})\otimes\mathscr{I}_n(\vec{C})$. In $\mathscr{A}_{n,m}(\vec{C})/(\xi\text{-relation})\otimes\mathscr{I}_n(\vec{C})$, we have

---

[*] Since d' does not satisfy $d' \circ d' = 0$, the word "d'-cycle" does not in fact make sense. But we merely say that an element of $\mathscr{G}_{n,m}(\vec{C})$ is a d'-cycle if it lies in the kernel of d'.

$$\begin{aligned}\langle(1\otimes d')\widetilde{\gamma}_{n,m}\rangle_1 &= \sum_{\Gamma\in\mathscr{G}_{n,m}(\vec{C})}[\Gamma]\otimes\langle d'\Gamma\rangle_1\\ &= \sum_{i=1}^{m}\sum_{\substack{\Gamma'(p_i,q_i)_i\\ d(p_i)=d(q_i)}}\Big(d'^*\Gamma'(p_i,q_i)_i\Big)\otimes\langle\Gamma'(p_i,q_i)_i\rangle_1,\end{aligned}$$

where the second sum of the second line is taken over graphs of degree $(\eta_1,\ldots,\eta_m)$, $\eta_j=1$ $(j\neq i)$, $\eta_i=0$ such that $\beta(i)\in\mathrm{Se}(\Gamma)$ and where $d'^*:\mathscr{G}_{n,m-1,\vec{\eta}}(\vec{C})\to\mathscr{A}_{n,m}(\vec{C})$ is the linear map defined by

$$d'^*\Gamma'=\Big(\text{coefficient of }1\otimes\langle\Gamma'\rangle_1\text{ in }\sum_{\Gamma\in\mathscr{G}_{n,m}(\vec{C})}[\Gamma]\otimes\langle d'\Gamma\rangle_1\Big).$$

This is well-defined since there is a canonical reduced presentation in $\mathscr{I}_n(\vec{C})$. But $d'^*\Gamma'(p_i,q_i)_i$, $d(p_i)=d(q_i)$, is precisely the left hand side of the definition of the ξ-relation. □

2.5. **The chain complex of endomorphisms of an acyclic chain complex.**

For a finitely generated, based free acyclic chain complex (C_*,∂), $C_i=\mathbb{Z}^{P_i}$, we consider the $\mathbb{Z}$-module $\mathrm{End}_k(C_*)$ of endomorphisms $C_*\to C_{*+k}$ of homogeneous degree k. The boundary operator $\partial':\mathrm{End}_k(C_*)\to\mathrm{End}_{k-1}(C_*)$ is defined by

$$\partial' f=\partial\circ f+(-1)^{k+1}f\circ\partial.$$

Then the pair $(\mathrm{End}_*(C_*),\partial')$ forms a chain complex. By the canonical isomorphism

$$\mathrm{End}_k(C_*)\cong\bigoplus_{i\in\mathbb{Z}}C_{i+k}\otimes\mathrm{Hom}(C_i,\mathbb{Z})$$

and the Künneth theorem, one can show that the complex $(\mathrm{End}_*(C_*),\partial')$ is acyclic.

For example, $f\in\mathrm{End}_0(C_*)$ is a cycle iff $\partial' f=\partial f-f\partial=0$. In particular, $\mathrm{id}\in\mathrm{End}_0(C_*)$ is a cycle and hence is a boundary. So there exists $g\in\mathrm{End}_1(C_*)$ such that

$$\partial' g=\partial g+g\partial=\mathrm{id}.$$

We will call such an endomorphism g a *combinatorial propagator* for (C_*,∂). If two such endomorphisms g,g' are given, then the difference $g-g'$ is a ∂'-cycle, since $\partial'(g-g')=\mathrm{id}-\mathrm{id}=0$. So there exists $h\in\mathrm{End}_2(C_*)$ such that

$$\partial' h=\partial h-h\partial=g-g'.$$

2.6. **Combinatorial propagator and trace.**

We will construct a $(d+d')$-cycle in $\mathscr{H}_n(\vec{C})$ by using combinatorial propagators and the trace in [Fuk96]. Let Γ be a $\vec{C}$-labeled graph Γ with $p_i\in C_*^{(i)}$ on the input and $q_i\in C_*^{(i)}$ on the output. We put $k_i=d(p_i)-d(q_i)$. For a sequence $\vec{h}=(h^{(1)},\ldots,h^{(m)})$ of endomorphisms

$h^{(i)} \in \mathrm{End}_{k_i}(C^{(i)}_*)$, $i = 1, 2, \ldots, m$, we define the trace of Γ by

$\mathrm{Tr}_{\vec{h}}$ 

$= \prod_{i=1}^{a} h^{(i)}_{q_ip_i} \times \mathrm{Smooth}$ 

,

where Smooth is the smoothing operator:

Smooth  =  .

In particular, since each $(C^{(i)}_*, \partial^{(i)})$ is acyclic, there exists a combinatorial propagator, i.e., an endomorphism $g^{(i)} : C^{(i)}_* \to C^{(i)}_{*+1}$ of homogeneous degree 1 such that $\partial^{(i)}g^{(i)} + g^{(i)}\partial^{(i)} = \mathrm{id}$. As a special case of the above definition, the trace by combinatorial propagators $\vec{g} = (g^{(1)}, \ldots, g^{(m)})$ defines a linear map

$$\mathrm{Tr}_{\vec{g}} : \mathscr{G}_{n,m}(\vec{C}) \to \mathscr{G}_{n,m}.$$

Let $\mathscr{A}_{n,m} = \mathscr{G}_{n,m}/(*)$-relation, label change relation. Since $\mathrm{Tr}_{\vec{g}}((*)\text{-relation}) \subset (*)$-relation, the map $\mathrm{Tr}_{\vec{g}}$ descends to a well-defined map

$$\overline{\mathrm{Tr}}_{\vec{g}} : \mathscr{A}_{n,m}(\vec{C}) \to \mathscr{A}_{n,m}.$$

Lemma 2.2. $\overline{\mathrm{Tr}}_{\vec{g}}(\xi\text{-}relation) = 0$. *Hence* $\overline{\mathrm{Tr}}_{\vec{g}}$ *induces a well-defined map*

$$\widetilde{\mathrm{Tr}}_{\vec{g}} : \mathscr{A}_{n,m}(\vec{C})/(\xi\text{-}relation) \to \mathscr{A}_{n,m}.$$

Proof. For $p_i, q_i \in P^{(i)}_*$ with $d(p_i) = d(q_i)$, we have

$$\begin{aligned}
&\overline{\mathrm{Tr}}_{\vec{g}}\Big[\sum_{\substack{x_i \in P^{(i)}_* \\ d(x_i)=d(q_i)+1}} \partial^{(i)}_{x_ip_i} \cdot [\Gamma(x_i, q_i)_i] + \sum_{\substack{y_i \in P^{(i)}_* \\ d(y_i)=d(p_i)-1}} \partial^{(i)}_{q_iy_i} \cdot [\Gamma(p_i, y_i)_i] - \delta_{p_iq_i} \cdot [\Gamma(\emptyset, \emptyset)_i]\Big] \\
&=\overline{\mathrm{Tr}}_{\ldots,\partial^{(i)}g^{(i)}+g^{(i)}\partial^{(i)},\ldots}\Big[[\Gamma(p_i, q_i)_i]\Big] - \overline{\mathrm{Tr}}_{\vec{g}}\Big[\delta_{p_iq_i} \cdot [\Gamma(\emptyset, \emptyset)_i]\Big] \\
&=\overline{\mathrm{Tr}}_{\ldots,\mathrm{id},\ldots}\Big[[\Gamma(p_i, q_i)_i]\Big] - \overline{\mathrm{Tr}}_{\vec{g}}\Big[\delta_{p_iq_i} \cdot [\Gamma(\emptyset, \emptyset)_i]\Big] \\
&=\delta_{p_iq_i}\overline{\mathrm{Tr}}_{\ldots,\mathrm{id},\ldots}\Big[[\Gamma(p_i, q_i)_i] - [\Gamma(\emptyset, \emptyset)_i]\Big] = 0.
\end{aligned}$$

□

Lemma 2.3. *The following assertions hold.* ($\widetilde{\mathrm{Tr}}' = \widetilde{\mathrm{Tr}} \otimes 1$)

(1) $\langle (1 \otimes (d + d'))\widetilde{\mathrm{Tr}}'_{\vec{g}}(\widetilde{\gamma}_{n,m})\rangle = 0$.

(2) $\langle \widetilde{\mathrm{Tr}}'_{\vec{g}}(\widetilde{\gamma}_{n,m})\rangle \in \mathscr{A}_{n,m} \otimes \mathscr{H}_n(\vec{C})$ *does not depend on the choice of combinatorial propagators* $\vec{g}$.

Proof. The assertion (1) is immediate from Lemmas 2.1 and 2.2 and from the fact that $\widetilde{\mathrm{Tr}}'_{\vec{g}}$ commutes with d and d'. Now we shall prove (2). We prove the assertion for $\vec{g} = (g, g^{(2)}, \ldots, g^{(m)})$ and $\vec{g}' = (g', g^{(2)}, \ldots, g^{(m)})$, where g and g' are two combinatorial propagators for $(C_*^{(1)}, \partial^{(1)})$. As mentioned in §2.5, there exists an endomorphism $h \in \mathrm{End}_2(C_*^{(1)})$ such that $\partial^{(1)}h - h\partial^{(1)} = g - g'$. Then we have

$$
\begin{aligned}
&\langle \widetilde{\mathrm{Tr}}'_{g,\ldots}(\widetilde{\gamma}_{n,m})\rangle - \langle \widetilde{\mathrm{Tr}}'_{g',\ldots}(\widetilde{\gamma}_{n,m})\rangle \\
&= \widetilde{\mathrm{Tr}}'_{g-g',\ldots}\Big[\sum_{\substack{\Gamma(p_1,q_1)_1 \\ d(p_1)=d(q_1)+1}} [\Gamma(p_1,q_1)_1] \otimes \langle \Gamma(p_1,q_1)_1 \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{\partial^{(1)}h-h\partial^{(1)},\ldots}\Big[\sum_{\Gamma(p_1,q_1)_1} [\Gamma(p_1,q_1)_1] \otimes \langle \Gamma(p_1,q_1)_1 \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{h,\ldots}\Big[\sum_{\Gamma(p_1,q_1)_1} \sum_{\substack{x_1 \in P_*^{(1)} \\ d(x_1)=d(p_1)+1}} \partial^{(1)}_{x_1p_1} \cdot [\Gamma(x_1,q_1)_1] \otimes \langle \Gamma(p_1,q_1)_1 \rangle \\
&\qquad - \sum_{\Gamma(p_1,q_1)_1} \sum_{\substack{y_1 \in P_*^{(1)} \\ d(y_1)=d(q_1)-1}} \partial^{(1)}_{q_1y_1} \cdot [\Gamma(p_1,y_1)_1] \otimes \langle \Gamma(p_1,q_1)_1 \rangle \Big] \qquad (2.1) \\
&= \widetilde{\mathrm{Tr}}'_{h,\ldots}\Big[\sum_{\Gamma(p_1,q_1)_1} \sum_{x_1} [\Gamma(x_1,q_1)_1] \otimes \langle \partial^{(1)}(x_1,p_1) * \Gamma(p_1,q_1)_1 \rangle \\
&\qquad - \sum_{\Gamma(p_1,q_1)_1} \sum_{y_1} [\Gamma(p_1,y_1)_1] \otimes \langle \Gamma(p_1,q_1)_1 * \partial^{(1)}(q_1,y_1) \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{h,\ldots}\Big[\sum_{\substack{p_1',q_1' \in P_*^{(1)} \\ d(p_1')=d(q_1')+2}} \sum_{\Gamma(p_1',q_1')_1} [\Gamma(p_1',q_1')_1] \otimes \langle (\partial^{(1)} * \Gamma - \Gamma * \partial^{(1)})(p_1',q_1')_1 \rangle \Big].
\end{aligned}
$$

We show that this vanishes. We write $\vec{C}[j] = (C_*^{(1)}, \ldots, \widehat{C_*^{(j)}}, \ldots, C_*^{(m)})$ and $\vec{\eta}[j] = (\eta_1, \ldots, \widehat{\eta}_j, \ldots, \eta_m)$ for simplicity. If we define $d^* : \bigoplus_{j=1}^m \mathscr{G}_{n-1,m-1,\vec{\eta}[j]}(\vec{C}[j]) \to \mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ by

$$
d^*\Gamma = \Big(\text{coefficient of } 1 \otimes \Gamma \text{ in } \sum_{\Gamma' \in \mathscr{G}_{n,m,\vec{\eta}}(\vec{C})} \Gamma' \otimes d\Gamma' \Big),
$$

then $\mathrm{Im}\, d^* = (*)$-relation. By the relation $(d + d')\mathscr{G}_{n,*,\Sigma(1,\ldots,1)}(\vec{C})$ of $\mathscr{H}_n(\vec{C})$, the following identities hold in $\mathscr{A}_{n,m} \otimes \mathscr{H}_n(\vec{C})$

$$
\begin{aligned}
0 &= \widetilde{\mathrm{Tr}}'_{h,\dots}\Big[\sum_{\substack{p'_1,q'_1 \in P_*^{(1)} \\ d(p'_1)=d(q'_1)+2}} \sum_{\substack{\Gamma(p'_1,q'_1)_1 \in \\ \mathscr{G}_{n,m,(2,1,\dots,1)}(\vec{C})}} [\Gamma(p'_1,q'_1)_1] \otimes \langle (d+d')\Gamma(p'_1,q'_1)_1 \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{h,\dots}\Big[\sum_{\substack{p'_1,q'_1 \in P_*^{(1)} \\ d(p'_1)=d(q'_1)+2}} \sum_{\substack{\Gamma'(p'_1,q'_1)_1 \in \\ \bigoplus_j \mathscr{G}_{n-1,m-1,(2,1,\dots,1)[j]}(\vec{C}[j])}} [d^*\Gamma'(p'_1,q'_1)_1] \otimes \langle \Gamma'(p'_1,q'_1)_1 \rangle \\
&\qquad + \sum_{\substack{p'_1,q'_1 \in P_*^{(1)} \\ d(p'_1)=d(q'_1)+2}} \sum_{\substack{\Gamma(p'_1,q'_1)_1 \in \\ \mathscr{G}_{n,m,(2,1,\dots,1)}(\vec{C})}} [\Gamma(p'_1,q'_1)_1] \otimes \langle d'\Gamma(p'_1,q'_1)_1 \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{h,\dots}\Big[\sum_{\substack{p'_1,q'_1 \in P_*^{(1)} \\ d(p'_1)=d(q'_1)+2}} \sum_{\Gamma(p'_1,q'_1)_1} [\Gamma(p'_1,q'_1)_1] \otimes \langle (\partial^{(1)} * \Gamma - \Gamma * \partial^{(1)})(p'_1,q'_1)_1 \rangle \\
&\qquad + \sum_{\substack{p'_1,q'_1 \in P_*^{(1)} \\ d(p'_1)=d(q'_1)+2}} \sum_{i \neq 1} \sum_{\substack{p_i,q_i \in P_*^{(i)} \\ d(p_i)=d(q_i)+1}} \sum_{\Gamma(p'_1,q'_1)_1(p_i,q_i)_i} [\Gamma(p'_1,q'_1)_1(p_i,q_i)_i] \\
&\qquad\qquad \otimes \langle (\partial^{(i)} * \Gamma + \Gamma * \partial^{(i)})(p'_1,q'_1)_1(p_i,q_i)_i \rangle \\
&\qquad + \sum_{\substack{p'_1,q'_1 \in P_*^{(1)} \\ d(p'_1)=d(q'_1)+2}} \sum_{\Gamma(p'_1,q'_1)_1} \sum_{\substack{e \in \mathrm{Comp}(\Gamma(p'_1,q'_1)_1) \\ e \neq 1}} [\Gamma(p'_1,q'_1)_1] \otimes \langle d'_e\Gamma(p'_1,q'_1)_1 \rangle \Big].
\end{aligned}
$$

Here the fourth row agrees with (2.1) and the vanishing of the last three rows can be shown as follows: for each $p'_1, q'_1 \in P_*^{(1)}$ with $d(p'_1) = d(q'_1) + 2$, we have

$$
\begin{aligned}
&\widetilde{\mathrm{Tr}}'_{h,\dots,g^{(i)},\dots}\Big[\sum_{\substack{p_i,q_i \in P_*^{(i)} \\ d(p_i)=d(q_i)+1}} \sum_{\Gamma(p'_1,q'_1)_1(p_i,q_i)_i} [\Gamma(p'_1,q'_1)_1(p_i,q_i)_i] \\
&\qquad\qquad \otimes \langle (\partial^{(i)} * \Gamma + \Gamma * \partial^{(i)})(p'_1,q'_1)_1(p_i,q_i)_i \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{h,\dots,\partial^{(i)}g^{(i)}+g^{(i)}\partial^{(i)},\dots}\Big[\sum_{\substack{p'_i,q'_i \in P_*^{(i)} \\ d(p'_i)=d(q'_i)}} \sum_{\Gamma(p'_1,q'_1)_1(p'_i,q'_i)_i} [\Gamma(p'_1,q'_1)_1(p'_i,q'_i)_i] \\
&\qquad\qquad \otimes \langle \Gamma(p'_1,q'_1)_1(p'_i,q'_i)_i \rangle \Big] \\
&= \widetilde{\mathrm{Tr}}'_{h,\dots,\mathrm{id},\dots}\Big[\sum_{p'_i \in P_*^{(i)}} \sum_{\Gamma(p'_1,q'_1)_1(p'_i,p'_i)_i} [\Gamma(p'_1,q'_1)_1(p'_i,p'_i)_i] \otimes \langle \Gamma(p'_1,q'_1)_1(p'_i,p'_i)_i \rangle \Big].
\end{aligned}
$$

This cancels with the corresponding term in

$$
\widetilde{\mathrm{Tr}}'_{h,\dots}\Big[\sum_{\Gamma(p'_1,q'_1)_1} \sum_{\substack{e \in \mathrm{Comp}(\Gamma(p'_1,q'_1)_1) \\ e \neq 1}} [\Gamma(p'_1,q'_1)_1] \otimes \langle d'_e\Gamma(p'_1,q'_1)_1 \rangle \Big].
$$

□

3. Fukaya's moduli space of gradient flow graphs in a manifold and its compactification

3.1. **Fukaya's moduli space $\mathscr{M}_\Gamma$.** Suppose given a sequence $\vec{f} = (f_1, f_2, \ldots, f_m)$ of C^r Morse functions on a d-dimensional smooth homology sphere M for $r \gg 0$ and a Riemannian metric μ on M. Throughout this section we assume that d is odd. Suppose that (f_i, μ) is Morse–Smale for each i. Recall that a pair of Morse function and a metric is Morse–Smale if all the intersections of the descending manifolds and the ascending manifolds are transversal. Let $C^{(i)} = (C^{(i)} = \mathbb{Z}^{P^{(i)}}, \partial^{(i)})$ be the Morse complex associated to (f_i, μ), namely, $C^{(i)}$ is the free $\mathbb{Z}$-module generated by the (finite) set $P_*^{(i)}$ of critical points of f_i and $\partial^{(i)} : C_{k+1}^{(i)} \to C_k^{(i)}$ is defined by

$$\begin{aligned}
&\partial^{(i)} p = \textstyle\sum_{q \in P_k^{(i)}} \#\overline{\mathscr{M}(f_i; p, q)} \cdot q,\\
&\overline{\mathscr{M}(f_i; p, q)} = \mathscr{M}(f_i; p, q) \pitchfork L_p,\\
&\mathscr{M}(f_i; p, q) = \mathscr{D}_p(f_i) \pitchfork \mathscr{A}_q(f_i),
\end{aligned}$$

where L_p is a level surface that lies just below the level of p and $\overline{\mathscr{M}(f_i; p, q)}$ is an oriented 0-dimensional manifold. Take a pair of critical points $a_i \in P_d^{(i)}$ and $b_i \in P_0^{(i)}$ arbitrarily, and consider the quotient

$$\widetilde{C}^{(i)} = C^{(i)} / \langle a_i, b_i \rangle.$$

Then $\widetilde{C}^{(i)}$ together with the induced boundary $\widetilde{\partial}^{(i)}$ forms an acyclic chain complex. We put

$$\vec{C} = (\widetilde{C}^{(1)}, \ldots, \widetilde{C}^{(m)}).$$

For a $\vec{C}$-colored graph $\Gamma = (\Gamma, \alpha, \beta)$ of type $(n, m, (1, \ldots, 1))$ with $\ell = m$, we define the source and the target maps

$$\mathrm{Source} : \{1, 2, \ldots, m\} \to \{1, 2, \ldots, n\}, \quad \mathrm{Target} : \{1, 2, \ldots, m\} \to \{1, 2, \ldots, n\}$$

as $\mathrm{Source}(k) = \alpha^{-1}(\text{source of } \beta(k))$, $\mathrm{Target}(k) = \alpha^{-1}(\text{target of } \beta(k))$. We define the subsets

$$\begin{aligned}
&\mathrm{In}_i(\Gamma) = \{k_1^i, k_2^i, \ldots, k_{r_i}^i\}, \quad \mathrm{In}_i^\infty(\Gamma) = \{\bar{k}_1^i, \bar{k}_2^i, \ldots, \bar{k}_{\bar{r}_i}^i\},\\
&\mathrm{Out}_i(\Gamma) = \{\ell_1^i, \ell_2^i, \ldots, \ell_{s_i}^i\}, \quad \mathrm{Out}_i^\infty(\Gamma) = \{\bar{\ell}_1^i, \bar{\ell}_2^i, \ldots, \bar{\ell}_{\bar{s}_i}^i\}
\end{aligned}$$

of the set of labels $\{1, 2, \ldots, m\}$ of edges as the subsets consisting of labels of edges with

$$\begin{aligned}
k_j^i \in \mathrm{In}_i(\Gamma) &\Leftrightarrow \mathrm{Target}(k_j^i) = i \text{ and } \beta(k_j^i) \in \mathrm{Comp}(\Gamma),\\
\bar{k}_j^i \in \mathrm{In}_i^\infty(\Gamma) &\Leftrightarrow \mathrm{Target}(\bar{k}_j^i) = i \text{ and } \beta(\bar{k}_j^i) \in \mathrm{Se}(\Gamma),\\
\ell_j^i \in \mathrm{Out}_i(\Gamma) &\Leftrightarrow \mathrm{Source}(\ell_j^i) = i \text{ and } \beta(k_j^i) \in \mathrm{Comp}(\Gamma),\\
\bar{\ell}_j^i \in \mathrm{Out}_i^\infty(\Gamma) &\Leftrightarrow \mathrm{Source}(\bar{\ell}_j^i) = i \text{ and } \beta(\bar{k}_j^i) \in \mathrm{Se}(\Gamma).
\end{aligned}$$

For example, $\mathrm{In}_i(\Gamma)$ is the subset of labels of incoming compact edges at the i-th vertex and $\mathrm{In}_i^\infty(\Gamma)$ is the subset of labels of incoming separated edges at the i-th vertex.

Let $\Phi_f^t : M \to M$ be the 1-parameter group of diffeomorphisms associated to $-\mathrm{grad}\, f$ considered with respect to a metric μ. For $\vec{f} = (f_1, \ldots, f_m)$, let $\mathscr{M}_\Gamma(\vec{f}) = \mathscr{M}_\Gamma(\vec{f}; \mu)$ be the space of points

$$(x_1, \ldots, x_n) \in \mathrm{Conf}_n(M)$$

such that

(1) for all $1 \leq i \leq n$, $1 \leq j \leq r_i$, there exists $t_{k_j^i} \in (0, \infty)$ such that $\Phi_{f_{k_j^i}}^{t_{k_j^i}}(x_{\sigma(k_j^i)}) = x_i$,

(2) $\lim_{t\to-\infty} \Phi_{f_{\bar{k}_j^i}}^t(x_i) = p_{\bar{k}_j^i}$ for $1 \leq i \leq n$, $1 \leq j \leq \bar{r}_i$,

(3) $\lim_{t\to+\infty} \Phi_{f_{\bar{\ell}_j^i}}^t(x_i) = q_{\bar{\ell}_j^i}$ for $1 \leq i \leq n$, $1 \leq j \leq \bar{s}_i$.

Remark 3.1. Since $\Phi_f^t(p) = \Phi_f^0(p) = p$ for a critical point p of f, we allow for a point $(x_1, \ldots, x_n)$ of $\mathscr{M}_\Gamma(\vec{f})$ that x_i coincides with a critical point of f_j. We will see below that such a point is not a singular point of $\mathscr{M}_\Gamma(\vec{f})$.

3.2. **Transversality.** Let $\Gamma \in \mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ be a $\vec{C}$-colored graph with a inputs (and a outputs) and with only regular edges. For simplicity, we assume that the non-compact edges in $\mathrm{Se}(\Gamma)$ are labeled (via β) by $\{1, 2, \ldots, a\}$. Let (p_i, q_i) be the pair of basis vectors attached on the edge labeled i. We define a smooth map

$$\Phi_{\vec{f}} : \prod_{j=1}^{a} (\mathscr{D}_{p_j}(f_j) \times \mathscr{A}_{q_j}(f_j)) \times \mathrm{Conf}_n(M) \times \mathbb{R}_+^{m-a} \to M^{n+m+a}$$

by

$$\begin{aligned}\Phi_{\vec{f}}&(u_1, v_1, \ldots, u_a, v_a; x_1, \ldots, x_n; t_{a+1}, \ldots, t_m) \\ &= \prod_{i=1}^{n} (x_i, y_{k_1^i}, \ldots, y_{k_{r_i}^i}, u_{\bar{k}_1^i}, \ldots, u_{\bar{k}_{\bar{r}_i}^i}, v_{\bar{\ell}_1^i}, \ldots, v_{\bar{\ell}_{\bar{s}_i}^i}),\end{aligned}$$

where $y_k = \Phi_{f_k}^{t_k}(x_{\sigma(k)})$ and $\mathbb{R}_+ = [0, \infty)$. Let $\Delta \subset M^{n+m+a}$ be the subset consisting of all the points of the form

$$\prod_{i=1}^{n} (x_i, \overbrace{x_i, \ldots, x_i}^{r_i+\bar{r}_i+\bar{s}_i})$$

for $(x_1, \ldots, x_n) \in \mathrm{Conf}_n(M)$. Then $\mathscr{M}_\Gamma(\vec{f})$ is the image of $\Phi_{\vec{f}}^{-1}(\Delta)$ under the projection onto $\mathrm{Conf}_n(M)$ and the projection induces an embedding.

Example 3.2. We consider the following graph, for example.

$\Gamma =$



$\in \mathscr{G}_{4,6}(\vec{C})$.

The map $\Phi_{\vec{f}} : (\mathscr{D}_p(f_1) \times \mathscr{A}_q(f_1)) \times \mathrm{Conf}_4(M) \times \mathbb{R}_+^5 \to M^{11}$ is defined by

$$\begin{aligned}\Phi_{\vec{f}}&(u, v; x_1, x_2, x_3, x_4; t_2, t_3, t_4, t_5, t_6) \\ &= (x_1, v) \times (x_2, \Phi_{f_2}^{t_2}(x_1)) \times (x_3, \Phi_{f_4}^{t_4}(x_1), \Phi_{f_5}^{t_5}(x_2)) \times (x_4, \Phi_{f_3}^{t_3}(x_2), \Phi_{f_6}^{t_6}(x_3), u).\end{aligned}$$

Then $\mathscr{M}_\Gamma(\vec{f})$ is the image of $\Phi_{\vec{f}}^{-1}(\Delta)$ under the projection onto $\mathrm{Conf}_4(M)$, where

$$\Delta=\{(x_1,x_1)\times(x_2,x_2)\times(x_3,x_3,x_3)\times(x_4,x_4,x_4,x_4)\,;\,(x_1,x_2,x_3,x_4)\in\mathrm{Conf}_4(M)\}.$$

□

Let $\mathscr{U}_j$ be a small neighborhood of a Morse function in $C^r(M)$, so that the set of critical points does not change on $\mathscr{U}_j$. By considering $\Phi_{\vec{f}}$ for all $\vec{f}\in\prod_{j=1}^m\mathscr{U}_j$ for a fixed Riemannian metric μ on M, we get a smooth map

$$\Phi:\prod_{j=1}^{a}\bigcup_{f_j\in\mathscr{U}_j}(\mathscr{D}_{p_j}(f_j)\times\mathscr{A}_{q_j}(f_j))\times\mathrm{Conf}_n(M)\times\mathbb{R}_+^{m-a}\times\prod_{j=a+1}^{m}\mathscr{U}_j\to M^{n+m+a},$$

where we consider $\bigcup_{f_j\in\mathscr{U}_j}(\mathscr{D}_{p_j}(f_j)\times\mathscr{A}_{q_j}(f_j))$ as a subspace of $\mathscr{U}_j\times M^2$.

Lemma 3.3. *The smooth map Φ is transversal to Δ.*

Proof. We show that every normal vector $V\in T_{\boldsymbol{x}}M^{m+n+a}$ lies in the image of the differential $D\Phi:T\mathrm{Domain}(\Phi)\to TM^{m+n+a}$ on points of Δ. Now suppose that a point $\boldsymbol{x}=\prod_{i=1}^n(x_i,\ldots,x_i)\in\Delta$ lies in the image of Φ. We first prove the claim in the case where each x_i does not belong to $\Sigma(f_1)\cup\cdots\cup\Sigma(f_m)$. Suppose that there exist $x\in M$ and $t_\ell\in(0,\infty)$ such that

$$\Phi_{f_\ell}^{t_\ell}(x)=x_i$$

for some ℓ. Then there exists a local coordinate $\psi_U:\mathbb{R}^d\overset{\approx}{\to}U$ of a neighborhood U of x_i in M such that $-\mathrm{grad}_{\psi_U(q)}f_\ell=D\psi_U(v_1(q),\ldots,v_d(q))$, $q\in\mathbb{R}^d$, $v_j(q)=v_j(0)$ (image of a constant field). Here, we may assume without loss of generality that $(D\psi_U)^{-1}(V)$ agrees with $(0,\ldots,0,b)$ for some $b<0$.

Now for small constants $\varepsilon>0$, $\kappa>0$ and $\beta>0$, we define a C^∞-function $g_{\varepsilon,\kappa}:\mathbb{R}\to\mathbb{R}$ by

$$g_{\varepsilon,\kappa}(t)=\begin{cases}-\kappa e^{-\frac{\beta t^2}{\varepsilon^2-t^2}} & -\varepsilon<t<\varepsilon\\ 0 & \text{otherwise}\end{cases}$$

We define a C^∞-function $h_\ell:\mathbb{R}^d\to\mathbb{R}$ by

$$h_\ell(t_1,\ldots,t_d)=f_\ell\circ\psi_U(t_1,\ldots,t_d)+sg_{\varepsilon,\kappa}(t_d).$$

for a constant $s>0$. By this deformation, the negative gradient, after passing through the part $-\varepsilon<t_d<\varepsilon$, shifts by a multiple of $(0,\ldots,0,1)$. Therefore, for some ε,κ and $\lambda_0\in\mathbb{R}$, we have

$$\left.\frac{d\Phi_{f_\ell+sg_{\varepsilon,\kappa}}^{t_\ell+s\lambda_0}}{ds}\right|_{s=0}=V.$$

This shows that the differential of Φ is surjective at any point of $\Phi^{-1}(\boldsymbol{x})$, $\boldsymbol{x}\in\Delta$.

If x_i agrees with $p_\ell\in\Sigma(f_\ell)$ for some ℓ and if the ℓ-th edge of Γ is one of the following forms:



then by a suitable deformation of f_ℓ on a small neighborhood of p_ℓ the position of p_ℓ shifts in an arbitrary direction. This shows the transversality for $x_i=p_\ell$ case. □

Proposition 3.4 (Fukaya). *Suppose that* $\Gamma \in \mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ *has no bivalent vertex, i.e.,* $E(\Gamma) = \mathrm{Reg}(\Gamma)$*. For a generic choice of* $\vec{f}$*, the space* $\mathscr{M}_\Gamma(\vec{f})$ *is an oriented smooth manifold of dimension*

$$\dim \mathscr{M}_\Gamma(\vec{f}) = d(n-m) + \sum_{i=1}^{m} \eta_i.$$

Moreover, $\vec{f}$ *can be chosen so that this property is satisfied simultaneously for all graphs* Γ *in* $\mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ *for a fixed triple* $m, n, \vec{\eta}$*.*

Proof. It follows from Lemma 3.3 that $\Phi^{-1}(\Delta)$ is a (infinite dimensional) smooth submanifold of codimension $(n+m+a-n)d = (m+a)d$. Let $\pi : \Phi^{-1}(\Delta) \to \prod_{j=1}^m \mathscr{U}_j$ be the restriction of the projection. Since the projection

$$\prod_{j=1}^{a} \bigcup_{f_j \in \mathscr{U}_j} (\mathscr{D}_{p_j}(f_j) \times \mathscr{A}_{q_j}(f_j)) \times \mathrm{Conf}_n(M) \times \mathbb{R}_+^{m-a} \times \prod_{j=a+1}^{m} \mathscr{U}_j \to \prod_{j=1}^{m} \mathscr{U}_j$$

is a Fredholm map of index $\sum_{j=1}^a (d(p_j) + (d - d(q_j))) + nd + (m-a) = nd + ad + m - a + \sum_{j=1}^a \eta_j$, the projection π is Fredholm of index

$$nd + ad + m - a + \sum_{j=1}^{a} \eta_j - (m+a)d = d(n-m) + \sum_{j=1}^{m} \eta_j.$$

Hence for a regular value $\vec{f} \in \prod_{j=1}^m \mathscr{U}_j$ of π, the fiber is a smooth manifold of dimension $d(n-m) + \sum_{j=1}^m \eta_j$. By the Sard–Smale theorem, the set of regular values of π is dense. The second statement follows from the fact that there are only finitely many graphs in $\mathscr{G}_{m,n,\vec{\eta}}(\vec{C})$ for a fixed $m, n, \vec{\eta}$ and that finite intersection of dense subsets is dense too. ☐

3.3. **Compactification of the moduli space of gradient trajectories between two non-singular points.**

For a Morse function f and a metric μ on M, we define

$$\mathscr{M}_2(f) = \mathscr{M}_2(f;\mu) = \{(x,y) \in (M - \Sigma(f))^2; y = \Phi_f^t(x) \text{ for some } t \in [0,\infty)\}.$$

We shall construct a natural compactification of $\mathscr{M}_2(f)$.

3.3.1. *The idea for the compactification.* To explain the idea for the compactification of $\mathscr{M}_2(f)$, we introduce the notion of a *(gradient) trajectory with nodes* as a sequence of points $x_0, x_1, \ldots, x_r \in M$ such that there is a sequence $t_0, t_1, \ldots, t_{r-1} \in [0,\infty)$ with $x_{i+1} = \Phi_f^{t_i}(x_i)$ for each i. Then it is obvious that $x_r = \Phi_f^{t_0+\cdots+t_{r-1}}(x_0)$, or $(x_0, x_r) \in \mathscr{M}_2(f)$. So, to describe a nonsingular trajectory connecting x_0 and x_r, the intermediate nodes $x_1, \ldots, x_{r-1}$ are unnecessary. However, to describe a broken trajectory as a limit of non-singular trajectories, they are necessary. For example, suppose a 1-parameter family γ_s of nonsingular trajectories between two nodes, say $x_0(s)$ and $x_2(s)$, converges to a twice broken trajectory ξ, broken at two critical points p and q, as $s \to \infty$. See Figure 2. The positions of $x_0(\infty)$ and $x_2(\infty)$ are not enough to determine ξ uniquely, since there may be not only one possibility for the trajectory connecting p and q. If we consider intermediate point, say $x_1(s)$, on γ_s, that converges to a point $x_1(\infty)$ with $f(q) < f(x_1(\infty)) < f(p)$, then the position



FIGURE 2. Convergence to a broken trajectory

of $x_1(\infty)$ determines the trajectory between p and q uniquely, and thus the limit ξ is determined uniquely as well. If the limit is k-times broken trajectory, then it suffices to consider appropriate family of nodes $x_1(s), \ldots, x_{k-1}(s)$ to describe the limiting behavior.

This observation suggests that the moduli space should be defined as the space of "equivalence classes" of trajectories with nodes, where we say that two trajectories with nodes $(x_0, x_1, \ldots, x_r)$ and $(y_0, y_1, \ldots, y_{r'})$ respectively are equivalent if $x_0 = y_0$ and $x_r = y_{r'}$. Then it will turn out that the limiting behavior of the moduli space near the end is nonsingular, in the sense that the moduli space can be considered as the interior of a smooth manifold with corners. See Proposition 3.10 for precise statement.

3.3.2. *Moduli space of short trajectories.* From now on we shall construct a compactification of $\mathscr{M}_2(f)$. One can easily check the following lemma.

Lemma 3.5. *For any C^r Morse function $f : M \to \mathbb{R}$, there is an arbitrarily C^r-small deformation of f in $C^r(M)$ such that all the critical values of the resulting Morse function are distinct.*

So, in the following, we assume that f is chosen as in Lemma 3.5. Since a smooth function on M that is sufficiently close in $C^r(M)$ to a Morse function is again a Morse function, we may assume that f is Morse. Suppose that the singular set $\Sigma(f) = \{p_1, p_2, \ldots, p_N\}$ is ordered so that $f(p_k) < f(p_{k+1})$ for each $k \leq N-1$. We put $c_k = f(p_k)$. For a small number $\varepsilon > 0$, let $L_k = f^{-1}(c_k - \varepsilon)$, $L'_k = f^{-1}(c_k + \varepsilon)$ and $W_k = f^{-1}[c_k - \varepsilon, c_{k+1})$. Note that W_k contains exactly one critical point of f. (See Figure 3.) For a pair of subsets A, B of M, we write

$$\mathscr{M}_2(f; A, B) = \mathscr{M}_2(f) \cap (A \times B).$$

We choose a small open cover $\mathscr{U} = \{U_\lambda\}$ of W_k. For each λ, we shall describe $\mathscr{M}_2(f; U_\lambda, L_k)$ and its compactification. There are three cases for λ:

(1) $U_\lambda \cap \mathscr{A}_{p_k}(f) = \emptyset$,
(2) $U_\lambda \cap \{p_k\} \neq \emptyset$ (and thus $U_\lambda \cap \mathscr{A}_{p_k}(f) \neq \emptyset$),
(3) $U_\lambda \cap \{p_k\} = \emptyset$ and $U_\lambda \cap \mathscr{A}_{p_k}(f) \neq \emptyset$.



FIGURE 3.

Case (1): In this case, the negative gradient trajectory starting from any point x of U_λ intersects L_k. We denote the intersection point on L_k by $\mu(x)$. Then the mapping $x \mapsto (x, \mu(x))$ defines a smooth embedding $\bar{\mu} : U_\lambda \to U_\lambda \times L_k$ and we have $\mathscr{M}_2(f; U_\lambda, L_k) = \operatorname{Im} \bar{\mu}$. One can see that there are no ends of $\operatorname{Im} \bar{\mu}$ in the image of the interior of U_λ, so it is not necessary to compactify the moduli space here, namely, we define

$$\overline{\mathscr{M}}_2(f; U_\lambda, L_k) = \mathscr{M}_2(f; U_\lambda, L_k).$$

Case (2): In this case, $\mathscr{M}_2(f; U_\lambda, L_k)$ is a submanifold of $(U_\lambda \setminus \mathscr{A}_{p_k}(f)) \times L_k$ diffeomorphic to $U_\lambda \setminus \mathscr{A}_{p_k}(f)$. By Morse's lemma, there is a local coordinate of a neighborhood M_{p_k} of p_k on which f agrees with

$$h(x) = c_k - x_1^2 - \cdots - x_i^2 + x_{i+1}^2 + \cdots + x_d^2.$$

One can choose a metric on M so that the metric on M_{p_k} corresponds to the standard metric on $\mathbb{R}^d$. We choose ε and U_λ so small that a neighborhood of $\mathscr{D}_{p_k}(f) \cap L_k$ in L_k is included in M_{p_k} and that $U_\lambda \subset M_{p_k}$. By the local coordinate of M_{p_k}, we have the following local description.

$$\mathscr{A}_{p_k}(h) = \{(x_1, \ldots, x_d) \in M_{p_k} \,;\, x_1 = \cdots = x_i = 0\},$$
$$\mathscr{D}_{p_k}(h) = \{(x_1, \ldots, x_d) \in M_{p_k} \,;\, x_{i+1} = \cdots = x_d = 0\}.$$

To simplify the argument, we replace L_k with

$$\overline{L}_k = \{(x_1, \ldots, x_d) \in M_{p_k} \,;\, x_1^2 + \cdots + x_i^2 = \varepsilon\} \subset M_{p_k},$$

which is tangent to L_k along $L_k \cap \mathscr{D}_{p_k}(f)$. This replacement does not cause any trouble since we are interested only in the local form of the moduli space of trajectories on a neighborhood of $\mathscr{A}_{p_k}(h) \cup \mathscr{D}_{p_k}(h)$. Under the local coordinate of M_{p_k}, $U_\lambda \setminus \mathscr{A}_{p_k}(h)$ can be considered as a subset of $(\mathbb{R}^i \setminus \{0\}) \times \mathbb{R}^{d-i}$, and similarly $\overline{L}_k = S^{i-1}_{\sqrt{\varepsilon}} \times \mathbb{R}^{d-i}$, where $S^{i-1}_{\sqrt{\varepsilon}} = \{(x_1, \ldots, x_i) \in \mathbb{R}^i ; x_1^2 + \cdots + x_i^2 = \varepsilon\}$.

By solving the differential equation

$$\frac{d}{dt}\gamma(t) = -\operatorname{grad}_{\gamma(t)} h,$$

we obtain

(3.1)

$$\begin{aligned}\mathscr{M}_2(f;U_\lambda,\overline{L}_k)=\Big\{(x_1,\ldots,x_d)\times\Big(\frac{\sqrt{\varepsilon}}{r}x_1,\ldots,\frac{\sqrt{\varepsilon}}{r}x_i,\frac{r}{\sqrt{\varepsilon}}x_{i+1},\ldots,\frac{r}{\sqrt{\varepsilon}}x_d\Big);\\ (x_1,\ldots,x_d)\in U_\lambda\setminus\mathscr{A}_{p_k}(h)\Big\},\end{aligned}$$

where $r=\sqrt{x_1^2+\cdots+x_i^2}$. Namely, $\mathscr{M}_2(f;U_\lambda,\overline{L}_k)$ agrees with the image of the restriction of an embedding $(\mathbb{R}^i\setminus\{0\})\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times(S^{i-1}_{\sqrt{\varepsilon}}\times\mathbb{R}^{d-i})$ given by the formula in (3.1) to $U_\lambda\setminus\mathscr{A}_{p_k}(h)$.

Now we shall construct a compactification of $\mathscr{M}_2(f;U_\lambda,\overline{L}_k)$ as the compactification of the embedding considered above. Let $\varphi_0:\mathbb{R}^i\setminus\{0\}\to\mathbb{R}^i\times S^{i-1}_{\sqrt{\varepsilon}}$ be the embedding defined by

$$\varphi_0(x_1,\ldots,x_i)=\Big(x_1,\ldots,x_i;\frac{\sqrt{\varepsilon}}{r}x_1,\ldots,\frac{\sqrt{\varepsilon}}{r}x_i\Big),$$

where $r=\sqrt{x_1^2+\cdots+x_i^2}$. The image of φ_0 has a canonical compactification as follows. Let $\widetilde{\gamma}^1(\mathbb{R}^i)$ denote the total space of the tautological oriented half-line ($[0,\infty)$) bundle over the oriented Grassmannian $\widetilde{G}_1(\mathbb{R}^i)\cong S^{i-1}_{\sqrt{\varepsilon}}$. Let $\pi:\widetilde{\gamma}^1(\mathbb{R}^i)\to\mathbb{R}^d$ be the map defined by $\pi=\mathrm{pr}_1\circ\varphi$ in the following diagram:

$$\begin{tikzcd}\widetilde{\gamma}^1(\mathbb{R}^i) \arrow[r, "\varphi"] \arrow[dr, "\pi"'] & \mathbb{R}^i\times S^{i-1}_{\sqrt{\varepsilon}} \arrow[r, "\mathrm{pr}_2"] \arrow[d, "\mathrm{pr}_1"] & S^{i-1}_{\sqrt{\varepsilon}} \\ & \mathbb{R}^i & \end{tikzcd}$$

where $\varphi:\widetilde{\gamma}^1(\mathbb{R}^i)\to\mathbb{R}^i\times S^{i-1}_{\sqrt{\varepsilon}}$ is the embedding, which maps a point u on a half-line $\ell\subset\mathbb{R}^i$ (starting at the origin) to the pair $(u,\ell\cap S^{i-1}_{\sqrt{\varepsilon}})$. Here, $\pi^{-1}(0)=\partial\widetilde{\gamma}^1(\mathbb{R}^i)$ is the image of the zero section of the tautological bundle $\mathrm{pr}_2\circ\varphi:\widetilde{\gamma}^1(\mathbb{R}^i)\to S^{i-1}_{\sqrt{\varepsilon}}$, that is diffeomorphic to S^{i-1}. Moreover, the restriction of π to the complement of $\pi^{-1}(0)$ is a diffeomorphism onto $\mathbb{R}^i\setminus\{0\}$ and the restriction of $\varphi\circ\pi^{-1}$ to $\mathbb{R}^i\setminus\{0\}$ agrees with φ_0. So $\mathrm{Im}\,\varphi$ is a compactification of $\mathrm{Im}\,\varphi_0$ to a smooth submanifold with boundary. (In fact, $\widetilde{\gamma}^1(\mathbb{R}^i)$ is the blow-up of 0 in $\mathbb{R}^i$.)

The compactification of $\mathscr{M}_2(f;U_\lambda,\overline{L}_k)$ is obtained from φ, as follows. Let $\varphi_0':(\mathbb{R}^i\setminus\{0\})\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times S^{i-1}_{\sqrt{\varepsilon}}$ be the embedding defined by

$$\begin{aligned}\varphi_0'(x_1,\ldots,x_d)&=(x_1,\ldots,x_d)\times\mathrm{pr}_2\circ\varphi_0(x_1,\ldots,x_i)\\&=(x_1,\ldots,x_d)\times\Big(\frac{\sqrt{\varepsilon}}{r}x_1,\ldots,\frac{\sqrt{\varepsilon}}{r}x_i\Big)\end{aligned}$$

and let $\varphi':\widetilde{\gamma}^1(\mathbb{R}^i)\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times S^{i-1}_{\sqrt{\varepsilon}}$ be its compactification:

$$\varphi'(u,\ell,x_{i+1},\ldots,x_d)=\pi(u,\ell)\times(x_{i+1},\ldots,x_d)\times\mathrm{pr}_2\circ\varphi(u,\ell).$$

This is just the direct product of φ with the identity on $\mathbb{R}^{d-i}$. Let $\varphi_0'':(\mathbb{R}^i\setminus\{0\})\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times(S^{i-1}_{\sqrt{\varepsilon}}\times\mathbb{R}^{d-i})$ be the embedding defined by

$$\varphi_0''(x_1,\ldots,x_d)=\varphi_0'(x_1,\ldots,x_d)\times\Big(\frac{r}{\sqrt{\varepsilon}}x_{i+1},\ldots,\frac{r}{\sqrt{\varepsilon}}x_d\Big)$$

and let $\varphi'' : \widetilde{\gamma}^1(\mathbb{R}^i) \times \mathbb{R}^{d-i} \to \mathbb{R}^d \times (S^{i-1}_{\sqrt{\varepsilon}} \times \mathbb{R}^{d-i})$ be the embedding defined by

$$\varphi''(u, \ell, x_{i+1}, \dots, x_d) = \varphi'(u, \ell, x_1, \dots, x_d) \times \Big(\frac{r}{\sqrt{\varepsilon}} x_{i+1}, \dots, \frac{r}{\sqrt{\varepsilon}} x_d \Big). \tag{3.2}$$

Now the following lemma is evident.

Lemma 3.6. *The following assertions hold.*

1. $\mathscr{M}_2(f; U_\lambda, \overline{L}_k) = \operatorname{Im} \varphi''_0 \cap ((U_\lambda \setminus \mathscr{A}_{p_k}(h)) \times \overline{L}_k)$.
2. $\operatorname{Im} \varphi'' = \overline{\operatorname{Im} \varphi''_0}$ *in* $\mathbb{R}^d \times (S^{i-1}_{\sqrt{\varepsilon}} \times \mathbb{R}^{d-i})$.
3. $\operatorname{Im} \varphi''$ *is a smooth submanifold with boundary.*
4. $\partial \operatorname{Im} \varphi'' = \Big\{ (0, \dots, 0, x_{i+1}, \dots, x_d) \times (x'_1, \dots, x'_i; 0, \dots, 0) \in \mathbb{R}^d \times (S^{i-1}_{\sqrt{\varepsilon}} \times \mathbb{R}^{d-i}) \Big\}$. *Hence* $\partial \operatorname{Im} \varphi'' \cap (U_\lambda \times \overline{L}_k) = (U_\lambda \cap \mathscr{A}_{p_k}(h)) \times (\overline{L}_k \cap \mathscr{D}_{p_k}(h))$. *Namely,* $\partial \operatorname{Im} \varphi'' \cap (U_\lambda \times \overline{L}_k)$ *is the space of once broken trajectories.*

We define

$$\overline{\mathscr{M}}_2(f; U_\lambda, \overline{L}_k) = \operatorname{Im} \varphi'' \cap (U_\lambda \times \overline{L}_k).$$

If U_λ is small enough, then the image of U_λ under the negative gradient flow is included in a small neighborhood of $\overline{L}_k \cap \mathscr{D}_{p_k}(h)$ in $\overline{L}_k$. Since $\overline{L}_k$ is tangent to L_k along $L_k \cap \mathscr{D}_{p_k}(f)$, the negative gradient flow gives a canonical embedding

$$\iota : \overline{\mathscr{M}}_2(f; U_\lambda, \overline{L}_k) \to U_\lambda \times L_k,$$

that induces a diffeomorphism $\operatorname{Int} \overline{\mathscr{M}}_2(f; U_\lambda, \overline{L}_k) \cong \mathscr{M}_2(f; U_\lambda, L_k)$. We define

$$\overline{\mathscr{M}}_2(f; U_\lambda, L_k) = \operatorname{Im} \iota.$$

Case (3): In this case, the negative gradient flow gives an isotopy

$$\Phi^t_f : U_\lambda \to M, \quad t \in [0, R]$$

for every $R > 0$, such that $\Phi^0_f = \mathrm{id}_{U_\lambda}$. Put $U'_\lambda = \Phi^R_f(U_\lambda)$. If U_λ is small enough, then for some choice of R, the open set U'_λ is included in M_{p_k} (a neighborhood of Morse's lemma). Then the image of the compactification of $\mathscr{M}_2(f; U'_\lambda, L_k)$ under the diffeomorphism $(\Phi^R_f)^{-1} \times \mathrm{id}_{L_k} : U'_\lambda \times L_k \to U_\lambda \times L_k$ gives a compactification $\overline{\mathscr{M}}_2(f; U_\lambda, L_k)$ of $\mathscr{M}_2(f; U_\lambda, L_k)$.

Now we define

$$\overline{\mathscr{M}}_2(f; W_k, L_k) = \bigcup_{U_\lambda \in \mathscr{U}} \overline{\mathscr{M}}_2(f; U_\lambda, L_k),$$

where $\overline{\mathscr{M}}_2(f; U_\lambda, L_k)$ and $\overline{\mathscr{M}}_2(f; U_\mu, L_k)$ are glued along $\overline{\mathscr{M}}_2(f; U_\lambda \cap U_\mu, L_k)$ through the inclusion induced maps. The compactifications $\overline{\mathscr{M}}_2(f; L_{k+1}, L_k)$, $\overline{\mathscr{M}}_2(f; L_{k+1}, W_k)$, $\overline{\mathscr{M}}_2(f; W_k, W_k)$ are similarly defined. Lemma 3.6 and its analogues imply the following proposition.

Proposition 3.7. $\overline{\mathscr{M}}_2(f; W_k, L_k)$ *(resp.* $\overline{\mathscr{M}}_2(f; L_{k+1}, L_k)$, $\overline{\mathscr{M}}_2(f; L_{k+1}, W_k)$*) is a smooth submanifold of* $W_k \times L_k$ *(resp.* $L_{k+1} \times L_k$, $L_{k+1} \times W_k$*) with corners, with*

$$\partial \overline{\mathscr{M}}_2(f; W_k, L_k) = \Big[(W_k \cap \mathscr{A}_{p_k}(f)) \times (L_k \cap \mathscr{D}_{p_k}(f)) \Big] \coprod \Delta_{L_k}$$

(resp. $(L_{k+1}\cap\mathscr{A}_{p_k}(f))\times(L_k\cap\mathscr{D}_{p_k}(f))$, $(L_{k+1}\cap\mathscr{A}_{p_k}(f))\times(W_k\cap\mathscr{D}_{p_k}(f))\coprod\Delta_{L_{k+1}}$*). The restriction of* $\overline{\mathscr{M}}_2(f;W_k,W_k)$ *on the complement of* Δ_{W_k} *is a smooth submanifold of* $W_k\times W_k$ *with corners, with*

$$\partial\overline{\mathscr{M}}_2(f;W_k,W_k)=\Big[(W_k\cap\mathscr{A}_{p_k}(f))\times(W_k\cap\mathscr{D}_{p_k}(f))\Big]\cup_{(p_k,p_k)}\Delta_{W_k}.$$

3.3.3. *Moduli space of long trajectories.* We will construct a compactification of $\mathscr{M}_2(f;W_k,W_j)$, $k>j$, by patching together the various compactifications obtained above. First, we construct a compactification of $\mathscr{M}_2(f;W_k,L_{k-1})$. The moduli space $\mathscr{M}_2(f;W_k,L_{k-1})$ can be identified with the intersection of $\mathscr{M}_2(f;W_k,L_k)\times\mathscr{M}_2(f;L_k,L_{k-1})\subset W_k\times L_k\times L_k\times L_{k-1}$ with $\widehat{\Delta}_{L_k}=\{(x,z_k,z_k,z_{k-1});x\in W_k,z_k\in L_k,z_{k-1}\in L_{k-1}\}$. It is easy to see that the intersection is transversal. Moreover, the transversality extends to the compactifications for a generic (f,μ), as in the following lemma.

Lemma 3.8. $\overline{\mathscr{M}}_2(f;W_k,L_k)\times\overline{\mathscr{M}}_2(f;L_k,L_{k-1})\subset W_k\times L_k\times L_k\times L_{k-1}$ *is transversal to* $\widehat{\Delta}_{L_k}$ *if* (f,μ) *satisfies the Morse–Smale condition.*

Proof. Let x be a point on the intersection. The orthogonal complement of $T_x(W_k\times\Delta_{L_k}\times L_{k-1})\subset T_x(W_k\times L_k\times L_k\times L_{k-1})$ is $N_x\Delta_{L_k}$, where $N_x\Delta_{L_k}\subset T_x(L_k\times L_k)$ is the orthogonal complement of $T_x\Delta_{L_k}\subset T_x(L_k\times L_k)$. So it suffices to show that $T_x(\overline{\mathscr{M}}_2(f;W_k,L_k)\times\overline{\mathscr{M}}_2(f;L_k,L_{k-1}))$ projects surjectively onto $N_x\Delta_{L_k}$. But this follows from the Morse–Smale condition. □

For a Morse–Smale pair (f,μ), we define

$$\overline{\mathscr{M}}_2(f;W_k,L_{k-1})=\mathrm{pr}\Big[\overline{\mathscr{M}}_2(f;W_k,L_k)\times\overline{\mathscr{M}}_2(f;L_k,L_{k-1})\cap\widehat{\Delta}_{L_k}\Big],$$

where $\mathrm{pr}:\widehat{\Delta}_{L_k}\to W_k\times L_k\times L_{k-1}$ is the projection $(x,z_k,z_k,z_{k-1})\mapsto(x,z_k,z_{k-1})$.

Lemma 3.9. *If* (f,μ) *is Morse–Smale,* $\overline{\mathscr{M}}_2(f;W_k,L_{k-1})$ *is a smooth manifold with corners, whose boundary is given by*

$$\begin{aligned}&\mathrm{pr}\Big[\partial\overline{\mathscr{M}}_2(f;W_k,L_k)\times\mathscr{M}_2(f;L_k,L_{k-1})\cap\widehat{\Delta}_{L_k}\Big]\\ \cup\,&\mathrm{pr}\Big[\mathscr{M}_2(f;W_k,L_k)\times\partial\overline{\mathscr{M}}_2(f;L_k,L_{k-1})\cap\widehat{\Delta}_{L_k}\Big]\\ \cup\,&\mathrm{pr}\Big[\partial\overline{\mathscr{M}}_2(f;W_k,L_k)\times\partial\overline{\mathscr{M}}_2(f;L_k,L_{k-1})\cap\widehat{\Delta}_{L_k}\Big].\end{aligned}$$

The interior of $\overline{\mathscr{M}}_2(f;W_k,L_{k-1})$ *is canonically diffeomorphic to* $\mathscr{M}_2(f;W_k,L_{k-1})$.

By repeating similar extensions inductively, we will obtain a compactification $\overline{\mathscr{M}}_2(f;W_k,W_j)$ of $\mathscr{M}_2(f;W_k,W_j)$ as a submanifold of $W_k\times L_k\times L_{k-1}\times\cdots\times L_{j+1}\times W_j$ with corners whose boundary is given by the subspace of broken trajectories.

3.3.4. *Moduli space of general trajectories.* We consider the gluing of $\overline{\mathscr{M}}_2(f;W_k,W_j)$ and $\overline{\mathscr{M}}_2(f;W_{k-1},W_j)$ to construct $\overline{\mathscr{M}}_2(f;W_k\cup W_{k-1},W_j)$ as follows. We consider $\overline{\mathscr{M}}_2(f;W_k\cap W_{k-1},W_j)$ as a submanifold of $\overline{\mathscr{M}}_2(f;W_k,W_j)$ and let $i_k:\overline{\mathscr{M}}_2(f;W_k\cap W_{k-1},W_j)\to\overline{\mathscr{M}}_2(f;W_k,W_j)$ be the inclusion. Let $j_k:\overline{\mathscr{M}}_2(f;W_k\cap W_{k-1},W_j)\to\overline{\mathscr{M}}_2(f;W_{k-1},W_j)$ be the codimension 0 embedding given by

$$j_k(x,z_k,z_{k-1},z_{k-2},\ldots,z_j)=(x,z_k,z_{k-2},z_{k-3},\ldots,z_j).$$

Namely, j_k forgets the node z_{k-1}. This is an embedding since $W_k \cap W_{k-1}$ does not include critical point of f. Then we define

$$\overline{\mathscr{M}}_2(f;W_k \cup W_{k-1}, W_j) = \overline{\mathscr{M}}_2(f;W_k,W_j) \cup_{\overline{\mathscr{M}}_2(f;W_k\cap W_{k-1},W_j)} \overline{\mathscr{M}}_2(f;W_{k-1},W_j),$$

where the gluing map is given by $j_k \circ i_k^{-1}$.

Extending in this way, we will finally obtain the moduli space $\overline{\mathscr{M}}_2(f;M,W_j)$. Moreover, a similar extension is valid also for the second factor W_j and we will finally get $\overline{\mathscr{M}}_2(f,M,M)$. We will write

$$\overline{\mathscr{M}}_2(f) = \overline{\mathscr{M}}_2(f;M,M).$$

Let $\overline{\Delta}_M$ be the closure of the lift of the diagonal $\Delta_{M-\Sigma(f)}$ in $\overline{\mathscr{M}}_2(f)$. The following proposition is evident.

Proposition 3.10. *Let (f,μ) be a Morse–Smale pair that is ordered as in Lemma 3.5. Then $\overline{\mathscr{M}}_2(f)$ is a smooth compact manifold with corners on the complement of $\overline{\Delta}_M$, whose boundary is given by the space of broken trajectories. The interior of $\overline{\mathscr{M}}_2(f)$ is canonically diffeomorphic to the interior of $\mathscr{M}_2(f)$.*

Remark 3.11. The compactification $\overline{\mathscr{M}}_2(f)$ is not a submanifold of $M\times M$ whereas $\mathscr{M}_2(f)$ is a submanifold of $M\times M$. There is a canonical continuous extension $\bar{\iota} : \overline{\mathscr{M}}_2(f) \to M\times M$ of the inclusion $\iota : \mathscr{M}_2(f) \to M \times M$, that assigns to each trajectory with nodes the pair of boundary nodes. The image of $\bar{\iota}$ is not a submanifold of $M\times M$ with corners since the dimension of some faces of the boundary decreases.

3.3.5. *Compactification of descending and ascending manifolds.* By Proposition 3.7, the subspace $\{p_k\}\times(L_k\cap\mathscr{D}_{p_k}(f))$ of $\overline{\mathscr{M}}_2(f;W_k,L_k)$ is a smooth submanifold of $\partial\overline{\mathscr{M}}_2(f;W_k,L_k)$. So, one could construct a compactification $\overline{\mathscr{M}}_2(f;\{p_k\},M)$ by a similar construction as above if one starts from $\{p_k\}\times(L_k\cap\mathscr{D}_{p_k}(f))$. We write

$$\overline{\mathscr{D}}_p(f) = \overline{\mathscr{M}}_2(f;\{p\},M).$$

One could also construct a compactification $\overline{\mathscr{A}}_p(f) = \overline{\mathscr{M}}_2(f;M,\{p\})$ similarly for some j. By a similar argument as given in the proof of Proposition 3.10, the following well-known proposition follows.

Proposition 3.12. *Let (f,μ) be a Morse–Smale pair that is ordered as in Lemma 3.5. Then $\mathscr{D}_p(f)$ (resp. $\mathscr{A}_p(f)$) has a natural compactification to a smooth manifold with corners $\overline{\mathscr{D}}_p(f)$ (resp. $\overline{\mathscr{A}}_p(f)$), whose codimension k stratum is*

$$\overline{\mathscr{D}}_p(f)_k = \bigcup_{\substack{q_1,\ldots,q_k\in\Sigma(f)\\ p,q_1,\ldots,q_k \text{ distinct}}} \overline{\mathscr{M}(f;p,q_1)}\times\overline{\mathscr{M}(f;q_1,q_2)}\times\cdots\times\overline{\mathscr{M}(f;q_{k-1},q_k)}\times\mathscr{D}_{q_k}(f),$$

$$\overline{\mathscr{A}}_p(f)_k = \bigcup_{\substack{q_1,\ldots,q_k\in\Sigma(f)\\ p,q_1,\ldots,q_k \text{ distinct}}} \mathscr{A}_{q_k}(f)\times\overline{\mathscr{M}(f;q_k,q_{k-1})}\times\cdots\times\overline{\mathscr{M}(f;q_2,q_1)}\times\overline{\mathscr{M}(f;q_1,p)}.$$

3.4. **Compactification of** $\mathrm{Conf}_n(M)$ **of Fulton–MacPherson.** For a closed d-dimensional manifold M, the configuration space $\mathrm{Conf}_n(M)$ is a smooth submanifold of M^n, that is the complement of the closed subset

$$\Sigma = \{(x_1,\ldots,x_n)\in M^n\,;\, x_i = x_j \text{ for some } i\neq j\}\subset M^n.$$

There is a natural filtration $\Sigma=\Sigma_n\supset\Sigma_{n-1}\supset\cdots\supset\Sigma_1\supset\Sigma_0$ with

$$\Sigma_j=\{(x_1,\ldots,x_n)\in M^n\,;\,\#\{x_1,\ldots,x_n\}=j+1\}.$$

The difference $\Sigma_{i+1}-\Sigma_i$ is a disjoint union of smooth submanifolds of $M^n-\Sigma_i$. This property allows one to iterate (real) blow-ups along the filtration from the deepest one: First, one can consider the blow-up $B\ell(M^n,\Sigma_0)$ along the d-dimensional submanifold Σ_0 of M^n. Recall that a blow-up replaces a submanifold with its normal sphere bundle. Since the closure of $\Sigma_1-\Sigma_0$ in $B\ell(M^n,\Sigma_0)$ is also a disjoint union of smooth submanifolds (with boundaries), one can apply another blow-up along it, and so on. After the blow-ups along all the strata of Σ of codimension ≥ 1, one obtains a smooth compact manifold with corners $\overline{\mathrm{Conf}}_n(M)$.

We will need a precise description of the boundary of $\overline{\mathrm{Conf}}_n(M)$ in the proof of invariance of $\widehat{Z}_{2k,3k}$, so we shall briefly recall it here. The space $\overline{\mathrm{Conf}}_n(M)$ has a natural stratification corresponding to bracketings of the n letters $1,2,\ldots,n$, e.g., ((137)(25))46 (see [FM]). Roughly speaking, a pair of brackets corresponds to a face created by one blow-up. For example, the face corresponding to ((137)(25))46 is obtained by a sequence of blow-ups corresponding to a sequence 1234567 → (12357)46 → ((137)(25))46.

The codimension one (boundary) strata of $\overline{\mathrm{Conf}}_n(M)$ correspond to bracketings of the form $(\cdots)\cdots$, with only one pair of brackets. The stratum $\partial_{\{1,\ldots,j\}}\overline{\mathrm{Conf}}_n(M)$ of $\partial\overline{\mathrm{Conf}}_n(M)$ corresponding to the bracketings $(12\cdots j)j{+}1\cdots n$ is the face created by the blow-up along the closure of the following submanifold of M^n:

$$\Delta_j=\{(x_1,\ldots,x_n)\in M^n\,;\,x_1=\cdots=x_j,\text{otherwise distinct}\}$$

in the result of the previous blow-ups. More precisely, $\partial_{\{1,\ldots,j\}}\overline{\mathrm{Conf}}_n(M)$ can be naturally identified with the blow-ups of the total space of the normal $S^{(j-1)d-1}$-bundle of $\Delta_j\subset M^n$ along the intersection with the closures of deeper diagonals that correspond to deeper bracketings. The fiber of the normal $S^{(j-1)d-1}$-bundle over a point $(x_j,\ldots,x_n)\in\Delta_j$ is

$$(\{(0,y_2,\ldots,y_j)\in(\mathbb{R}^d)^j\}-\{0\})/(\text{dilation})\cong S^{(j-1)d-1},$$

where the coordinate y_i corresponds to x_i-x_1 (where it makes sense) via a local framing of $T_{x_j}M$. The stratum $\partial_{\{1,\ldots,j\}}\overline{\mathrm{Conf}}_n(M)$ is a fiber bundle over Δ_j. We denote the fiber of $\partial_{\{1,\ldots,j\}}\overline{\mathrm{Conf}}_n(M)$ over a point of Δ_j by $\mathrm{Conf}_j^{\mathrm{local}}(\mathbb{R}^d)$. There is a canonical identification of $\mathrm{Conf}_j^{\mathrm{local}}(\mathbb{R}^d)$ with the subset of $\mathrm{Conf}_j(\mathbb{R}^d)$ consisting of configurations $(y_1,\ldots,y_j)$ such that $y_1=0$ and $\sum_{\ell=2}^j\|y_\ell\|^2=1$. We denote by $\overline{\mathrm{Conf}}_j^{\mathrm{local}}(\mathbb{R}^d)$ the closure of the image of the inclusion $\mathrm{Conf}_j^{\mathrm{local}}(\mathbb{R}^d)\hookrightarrow\mathrm{Conf}_j(M)\hookrightarrow\overline{\mathrm{Conf}}_j(\mathbb{R}^d)$. The base space Δ_j is naturally diffeomorphic to $\overline{\mathrm{Conf}}_{n-j+1}(M)$ and we denote by $\mathrm{pr}_j:\Delta_j\to M$ the projection $(x_j,\ldots,x_n)\mapsto x_j$. So $\partial_{\{1,\ldots,j\}}\overline{\mathrm{Conf}}_n(M)$ has the structure of the pullback of the associated $\overline{\mathrm{Conf}}_j^{\mathrm{local}}(\mathbb{R}^d)$-bundle of TM ($\overline{\mathrm{Conf}}_j^{\mathrm{local}}(\mathbb{R}^d)$ is an SO_d-space) pulled back by pr_j. The definition of $\partial_A\overline{\mathrm{Conf}}_n(M)$ for a subset $A\subset\{1,\ldots,n\}$ corresponding to the bracketing $(A)A^c$ is similar.

It will turn out that the faces of $\partial\overline{\mathrm{Conf}}_n(M)$ corresponding to coincidence of two points are special among the codimension one strata of $\overline{\mathrm{Conf}}_n(M)$. We denote by

$\partial^{\mathrm{pri}}\overline{\mathrm{Conf}}_n(\mathbb{R}^d)$ ("pri" for principal) the union of the faces corresponding to coincidence of two points and we denote by $\partial^{\mathrm{hi}}\overline{\mathrm{Conf}}_n(\mathbb{R}^d)$ ("hi" for hidden) the union of all the faces corresponding to coincidence of at least three points.

3.5. **Compactification of Fukaya's moduli space of gradient flow graphs in a manifold.** We shall construct a compactfication $\overline{\mathscr{M}}_\Gamma(\vec{f})$ of $\mathscr{M}_\Gamma(\vec{f})$. We shall first give another definition of $\mathscr{M}_\Gamma(\vec{f})$ using $\mathscr{M}_2(f)$. For simplicity, we again assume that the non-compact edges in $\mathrm{Se}(\Gamma)$ are labeled (via β) by $\{1, 2, \ldots, a\}$. We define an embedding

$$\Psi_{\vec{f}} : \prod_{j=1}^{a}(\mathscr{D}_{p_j}(f_j) \times \mathscr{A}_{q_j}(f_j)) \times \prod_{j=a+1}^{m} \mathscr{M}_2(f_j) \to M^{2m}$$

as follows. For simplicity, we only give a definition of $\Psi_{\vec{f}}$ for Γ of Example 3.2. In that case, we define

$$\Psi_{\vec{f}}(u, v; x_2, y_2; \ldots; x_6, y_6) = (v, x_2, x_4) \times (x_3, x_5, y_2) \times (x_6, y_4, y_5) \times (u, y_3, y_6).$$

Let C^Δ be the subspace of M^{12} defined by

$$\{(z_1, z_1, z_1) \times (z_2, z_2, z_2) \times (z_3, z_3, z_3) \times (z_4, z_4, z_4)\,;\ z_i \in M,\ z_i \neq z_j \text{ if } i \neq j\}.$$

Then there exists a canonical diffeomorphism $\varpi : C^\Delta \to \mathrm{Conf}_4(M)$ and one may see that

$$\varpi(\mathrm{Im}\,\Psi_{\vec{f}} \cap C^\Delta) = \mathscr{M}_\Gamma(\vec{f}).$$

We will construct a compactification of $\mathrm{Im}\,\Psi_{\vec{f}} \cap C^\Delta$. There is a straightforward extension

$$\overline{\Psi}_{\vec{f}} : \prod_{j=1}^{a}(\overline{\mathscr{D}}_p(f_j) \times \overline{\mathscr{A}}_q(f_j)) \times \prod_{j=a+1}^{m} \overline{\mathscr{M}}_2(f_j) \to M^{2m}$$

of $\Psi_{\vec{f}}$. For example, for Γ of Example 3.2,

$$\overline{\Psi}_{\vec{f}}(u, v; x_2, y_2; \ldots; x_6, y_6) = (\sigma(v), x_2, x_4) \times (x_3, x_5, y_2) \times (x_6, y_4, y_5) \times (\tau(u), y_3, y_6),$$

where $\tau : \overline{\mathscr{D}}_p(f_1) \to M$ (resp. $\sigma : \overline{\mathscr{A}}_q(f_1) \to M$) be the continuous map that assigns the terminal point (resp. source point) in a (possibly broken) trajectory, that extend the inclusions $\mathscr{D}_p(f_1) \hookrightarrow M$ (resp. $\mathscr{A}_q(f_1) \hookrightarrow M$). There is a canonical embedding $\gamma : \mathrm{Conf}_n(M) \to \overline{\mathrm{Conf}}_n(M)$, that induces a diffeomorphism onto the interior. Then we define

$$\overline{\mathscr{M}}_\Gamma(\vec{f}) = \mathrm{Closure}\big(\gamma\mathscr{M}_\Gamma(\vec{f})\big) = \mathrm{Closure}\big(\gamma \circ \varpi(\mathrm{Im}\,\overline{\Psi}_{\vec{f}} \cap C^\Delta)\big) \subset \overline{\mathrm{Conf}}_n(M).$$

The following proposition follows immediately.

Proposition 3.13. (1) $\mathrm{Int}\,\overline{\mathscr{M}}_\Gamma(\vec{f}) = \gamma\mathscr{M}_\Gamma(\vec{f})$. *The space* $\overline{\mathscr{M}}_\Gamma(\vec{f})$ *is compact.*

(2) *On a neighborhood* U *of a point on the strata of codimension* ≤ 1 *of* $\mathrm{Domain}(\overline{\Psi}_{\vec{f}})$, $\overline{\Psi}_{\vec{f}}$ *is transversal to* C^Δ. *Thus* $\overline{\mathscr{M}}_\Gamma(\vec{f}) \cap \overline{\Psi}_{\vec{f}}(U)$ *is a smooth submanifold with boundary.*

(3) $\partial(\overline{\mathscr{M}}_\Gamma(\vec{f}) \cap \overline{\Psi}_{\vec{f}}(U))$

$$= \Big[\gamma \circ \varpi(\overline{\Psi}_{\vec{f}}\partial\mathrm{Domain}(\overline{\Psi}_{\vec{f}}) \cap C^\Delta) \cup \partial\overline{\mathrm{Conf}}_n(M)\Big] \cap (\overline{\mathscr{M}}_\Gamma(\vec{f}) \cap \overline{\Psi}_{\vec{f}}(U)).$$

Namely, the codimension one stratum of $\overline{\mathscr{M}}_\Gamma(\vec{f})$ *consists of flows having*

(a) *one once broken edge, or*

(b) *one subgraph collapsed into a point.*

Remark 3.14. By Remark 3.11, $\overline{\mathscr{M}}_\Gamma(\vec{f})$ may not be a smooth submanifold of $\overline{\mathrm{Conf}}_n(M)$ on the image from the strata of codimension ≥ 2. To make this smooth submanifold, one needs to replace $\overline{\mathrm{Conf}}_n(M)$ with $\overline{\mathrm{Conf}}_n(M)\times M^r$ for some positive integer r. Then $\overline{\Psi}_{\vec{f}}$ induces an embedding of the smooth manifold $\overline{\Psi}_{\vec{f}}^{-1}(C^\Delta)$.

3.6. **Orientations of $\partial\overline{\mathscr{D}}_p(f)$ and $\partial\overline{\mathscr{A}}_q(f)$.** Let $f: M\to\mathbb{R}$ be a Morse function and μ be a metric on M that is Morse–Smale. We emphasize that we are assuming that $d=\dim M$ is odd. We shall describe the induced orientation on the codimension 1 stratum $\mathscr{F}_r\overline{\mathscr{D}}_p(f)$ of broken trajectories $\gamma(t)$ with $\lim_{t\to-\infty}\gamma(t)=p$ that is broken at a critical point r when (f,μ) is Morse–Smale. We consider $\mathscr{F}_r\overline{\mathscr{D}}_p(f)$ as a submanifold of $M\times M$. In the following we use the orientation convention of Appendix A.

3.6.1. *Orientation of* $\partial\overline{\mathscr{D}}_p(f)$. Let a be a point on $\mathscr{D}_r(f)$ that is close to r and choose a small neighborhood U_a of a in M, which is so small that U_a is included in a local model M_r of the Morse lemma around r. Let $c_r=f(r)$ and let $L'_r=f^{-1}(c_r+\varepsilon)$. Choose a local coordinate $(x_1,\ldots,x_d)\in\mathbb{R}^d$ on the Morse model M_r so that $0\in\mathbb{R}^d$ corresponds to r. If $f(x)-c_r$ agrees on M_r with the standard quadratic form $h(x)=-x_1^2-\cdots-x_i^2+x_{i+1}^2+\cdots+x_d^2$, then we put $\overline{L'_r}=\{x_{i+1}^2+\cdots+x_d^2=\varepsilon\}\subset\mathbb{R}^d$. We shall describe the orientation of $\partial\overline{\mathscr{D}}_p(f)_1\cap(\overline{L'_r}\times U_a)$ that is induced from the orientation of $\overline{\mathscr{D}}_p(f)$.

Suppose that $d(r)=i$, $d(p)=i+k+1$ (so that $\dim\overline{\mathscr{M}(f;p,r)}=k$). Let $\varphi''_0:\mathbb{R}^i\times(\mathbb{R}^{d-i}\setminus\{0\})\to(\mathbb{R}^i\times S^{d-i-1}_{\sqrt{\varepsilon}})\times\mathbb{R}^d\subset\mathbb{R}^d\times\mathbb{R}^d$ be the embedding defined by

$$\begin{aligned}&\varphi''_0(x_1,\ldots,x_i;x_{i+1},\ldots,x_d)\\&\quad=\Big(\frac{r}{\sqrt{\varepsilon}}x_1,\ldots,\frac{r}{\sqrt{\varepsilon}}x_i\Big)\times\Big(\frac{\sqrt{\varepsilon}}{r}x_{i+1},\ldots,\frac{\sqrt{\varepsilon}}{r}x_d\Big)\times(x_1,\ldots,x_d),\end{aligned}$$

where $r=\sqrt{x_{i+1}^2+\cdots+x_d^2}$ and let $\varphi'':\mathbb{R}^i\times\widetilde{\gamma}^1(\mathbb{R}^{d-i})\to(\mathbb{R}^i\times S^{d-i-1}_{\sqrt{\varepsilon}})\times\mathbb{R}^d$ be its compactification as above (see (3.2)). Then we have

$$\begin{aligned}\overline{\mathscr{M}}_2(f;\overline{L'_r},U_a)&=\mathrm{Im}\,\varphi''\cap(\overline{L'_r}\times U_a),\ \text{and}\\ \partial\,\mathrm{Im}\,\varphi''&=(\{0\}\times S^{d-i-1}_{\sqrt{\varepsilon}})\times(\mathbb{R}^i\times\{0\}).\end{aligned}$$

Note that the image of φ'' agrees with the following subspace

$$\begin{aligned}\Big\{&\Big(\frac{r}{\sqrt{\varepsilon}}x'_1,\ldots,\frac{r}{\sqrt{\varepsilon}}x'_i;x_{i+1},\ldots,x_d\Big)\times\Big(x'_1,\ldots,x'_i;\frac{r}{\sqrt{\varepsilon}}x_{i+1},\ldots,\frac{r}{\sqrt{\varepsilon}}x_d\Big);\\&\qquad(x_{i+1},\ldots,x_d)\in S^{d-i-1}_{\sqrt{\varepsilon}},\ (x'_1,\ldots,x'_i)\in\mathbb{R}^i,\ r\in[0,\infty)\Big\}.\end{aligned}$$

Let $b=(0,\ldots,0,\sqrt{\varepsilon})\in\overline{L'_r}$. By symmetry, we need only to consider the orientations on a neighborhood $\widetilde{U}_b$ of $\{b\}\times(\mathscr{D}_r(f)\cap U_a)\subset\overline{L'_r}\times U_a$. Moreover we assume for simplicity that $T_b(\mathscr{D}_p\cap\overline{L'_r})=\{(x_1,\ldots,x_{i+k},0,\ldots,0,\sqrt{\varepsilon})\}\subset\{(x_1,\ldots,x_{d-1},\sqrt{\varepsilon})\}=T_b\overline{L'_r}$. We put

$$\begin{aligned}o(\mathscr{D}_r(f))_x&=\alpha\,dx_1\wedge\cdots\wedge dx_i,\\ o(\mathscr{D}_p(f))_x&=\beta\,dx_1\wedge\cdots\wedge dx_{i+k}\wedge dx_d,\end{aligned}\tag{3.3}$$

for $x \in M_r$, $\alpha, \beta \in \{+1, -1\}$, where these make sense. This gives

$$\begin{aligned} o^*(\mathscr{A}_r(f))_x &= \alpha\, dx_1 \wedge \cdots \wedge dx_i, \\ o^*(\mathscr{D}_p(f))_x &= (-1)^{d-1-i-k} \beta\, dx_{i+k+1} \wedge \cdots \wedge dx_{d-1}. \end{aligned}$$

We also consider $\mathscr{D}_p(f) \cap U_a$ as a submanifold of $\mathbb{R}^d \times \mathbb{R}^d$ via the embedding φ_0''. The orientation of $\mathscr{D}_p(f) \cap U_a$, considered as a submanifold of $\mathbb{R}^d \times \mathbb{R}^d$, induced from (3.3) is

$$o(\mathscr{D}_p(f))_{(x,x')} = \beta\, dx_1' \wedge \cdots \wedge dx_i' \wedge dx_{i+1} \wedge \cdots \wedge dx_{i+k} \wedge dx_d.$$

Hence we obtain

$$\begin{aligned} &o^*(\mathscr{D}_p(f))_{(x,x')} \\ &= (-1)^i \beta\, dx_{i+1}' \wedge \cdots \wedge dx_d' \wedge dx_1 \wedge \cdots \wedge dx_i \wedge dx_{i+k+1} \wedge \cdots \wedge dx_{d-1}. \end{aligned}$$

This induces the following orientation on the boundary:

$$\begin{aligned} &(-1)^i(-1)^{2d-1}\beta\, dx_{i+1}' \wedge \cdots \wedge dx_d' \wedge dx_1 \wedge \cdots \wedge dx_i \wedge dx_{i+k+1} \wedge \cdots \wedge dx_d \\ =&(-1)^{i+1}\beta\, dx_{i+1}' \wedge \cdots \wedge dx_d' \wedge dx_1 \wedge \cdots \wedge dx_i \wedge dx_{i+k+1} \wedge \cdots \wedge dx_d. \end{aligned} \tag{3.4}$$

Note that the inward normal direction of $\partial\overline{\mathscr{M}}_2(f; \overline{L_r'}, U_a) \cap \widetilde{U}_b$ corresponds to the outward normal direction of $S^{d-i-1}_{\sqrt{\varepsilon}}$ (recall that $S^{d-i-1}_{\sqrt{\varepsilon}}$ corresponds to the boundary face of $\widetilde{\gamma}^1(\mathbb{R}^{d-i})$). So the dual of the inward vector field with respect to the standard metric is given by dx_d. Then apply (A.1).

Given these data, we compute $o^*(\overline{\mathscr{M}(f;p,r)})$ as follows.

$$\begin{aligned} o^*(\mathscr{M}_2(f;p,r))_x &= o^*(\mathscr{D}_p(f) \pitchfork \mathscr{A}_r(f))_x \\ &= o^*(\mathscr{D}_p(f))_x \wedge o^*(\mathscr{A}_r(f))_x \\ &= (-1)^{k(i+1)} \alpha\beta\, dx_1 \wedge \cdots \wedge dx_i \wedge dx_{i+k+1} \wedge \cdots \wedge dx_{d-1}. \end{aligned}$$

So we have

$$o^*(\overline{\mathscr{M}(f;p,r)})_x = (-1)^{k(i+1)} \alpha\beta\, dx_1 \wedge \cdots \wedge dx_i \wedge dx_{i+k+1} \wedge \cdots \wedge dx_{d-1} \wedge dx_d,$$

where dx_d corresponds to the dual of $\mathrm{grad}_x f$ with respect to the standard metric . Therefore,

$$\begin{aligned} &o^*(\overline{\mathscr{M}(f;p,r)})_x \wedge o^*(\mathscr{D}_r(f))_{x'} \\ &= (-1)^{i+1} \beta\, dx_{i+1}' \wedge \cdots \wedge dx_d' \wedge dx_1 \wedge \cdots \wedge dx_i \wedge dx_{i+k+1} \wedge \cdots \wedge dx_d. \end{aligned}$$

This agrees with the orientation of $\partial\overline{\mathscr{D}}_p(f)_1$ induced from $o^*(\mathscr{D}_p(f))_{(x,x')}$ given above in (3.4).

Lemma 3.15. *The normal orientation on $\partial\overline{\mathscr{D}}_p(f)_1 \cap (\overline{L_r'} \times U_a)$ induced from $o^*(\mathscr{D}_p(f))_{(x,x')}$ agrees with $o^*(\overline{\mathscr{M}(f;p,r)})_x \wedge o^*(\mathscr{D}_r(f) \times U_a)_{x'}$.*

3.6.2. *Orientation of* $\partial\overline{\mathscr{A}}_q(f)$. Suppose that $d(r)=i$ and $d(q)=i-k-1$ (so that $\dim\overline{\mathscr{M}(f;r,q)}=k$). The orientation of $\partial\overline{\mathscr{A}}_q(f)_1$ is described as follows. Let $\varphi_0'' : (\mathbb{R}^i\setminus\{0\})\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times(S^{i-1}_{\sqrt{\varepsilon}}\times\mathbb{R}^{d-i})$ be the embedding defined by

$$\begin{aligned}&\varphi_0''(x_1,\ldots,x_i;x_{i+1},\ldots,x_d)\\ &=(x_1,\ldots,x_d)\times\Big(\frac{\sqrt{\varepsilon}}{r}x_1,\ldots,\frac{\sqrt{\varepsilon}}{r}x_i\Big)\times\Big(\frac{r}{\sqrt{\varepsilon}}x_{i+1},\ldots,\frac{r}{\sqrt{\varepsilon}}x_d\Big),\end{aligned}\tag{3.5}$$

where $r=\sqrt{x_1^2+\cdots+x_i^2}$ and let $\varphi'' : \widetilde{\gamma}^1(\mathbb{R}^i)\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times(S^{i-1}_{\sqrt{\varepsilon}}\times\mathbb{R}^{d-i})$ be its compactification. Then we have

$$\begin{gathered}\overline{\mathscr{M}}_2(f;U_a,\overline{L}_r)=\operatorname{Im}\varphi''\cap(U_a\times\overline{L}_r)\text{ and}\\ \partial\operatorname{Im}\varphi''=(\{0\}\times\mathbb{R}^{d-i})\times(S^{i-1}_{\sqrt{\varepsilon}}\times\{0\}).\end{gathered}$$

Note that the image of φ'' agrees with the following subspace

$$\begin{aligned}\Big\{&\Big(\frac{r}{\sqrt{\varepsilon}}x_1',\ldots,\frac{r}{\sqrt{\varepsilon}}x_i';x_{i+1},\ldots,x_d\Big)\times\Big(x_1',\ldots,x_i';\frac{r}{\sqrt{\varepsilon}}x_{i+1},\ldots,\frac{r}{\sqrt{\varepsilon}}x_d\Big);\\ &\qquad(x_{i+1},\ldots,x_d)\in\mathbb{R}^{d-i},\ (x_1',\ldots,x_i')\in S^{i-1}_{\sqrt{\varepsilon}},\ r\in[0,\infty)\Big\}.\end{aligned}$$

Let $b=(\sqrt{\varepsilon},0,\ldots,0)\in\overline{L}_r$ and we consider the orientations on a neighborhood $\widetilde{U}_b$ of $(\mathscr{A}_r(f)\cap U_a)\times\{b\}\subset U_a\times\overline{L}_r$. Moreover we assume that $T_b(\mathscr{A}_q(f)\cap\overline{L}_r)=\{(\sqrt{\varepsilon},x_2,\ldots,x_{k+1},0,\ldots,0,x_{i+1},\ldots,x_d)\}$ for simplicity. We put

$$\begin{aligned}o(\mathscr{A}_r(f))_x&=\alpha\,dx_{i+1}\wedge\cdots\wedge dx_d,\\ o(\mathscr{A}_q(f))_x&=\beta\,dx_1\wedge\cdots\wedge dx_{k+1}\wedge dx_{i+1}\wedge\cdots\wedge dx_d,\end{aligned}\tag{3.6}$$

for $x\in M_r$, $\alpha,\beta\in\{+1,-1\}$, where these make sense. This gives

$$\begin{aligned}o^*(\mathscr{D}_r(f))_x&=\alpha\,dx_{i+1}\wedge\cdots\wedge dx_d,\\ o^*(\mathscr{A}_q(f))_x&=(-1)^{(i-k-1)(d-i)}\beta\,dx_{k+2}\wedge\cdots\wedge dx_i.\end{aligned}$$

We also consider $\mathscr{A}_q(f)\cap U_a$ as a submanifold of $\mathbb{R}^d\times\mathbb{R}^d$ via the embedding φ_0''. The orientation of $\mathscr{A}_q(f)\cap U_a$, considered as a submanifold of $\mathbb{R}^d\times\mathbb{R}^d$, induced from (3.6) is

$$o(\mathscr{A}_q(f))_{(x,x')}=\beta\,dx_1'\wedge\cdots\wedge dx_{k+1}'\wedge dx_{i+1}\wedge\cdots\wedge dx_d.$$

Hence we obtain

$$o^*(\mathscr{A}_q(f))_{(x,x')}=(-1)^{k+1}\beta\,dx_1\wedge\cdots\wedge dx_i\wedge dx_{k+2}'\wedge\cdots\wedge dx_d'.\tag{3.7}$$

This induces the following orientation on the boundary:

$$\begin{aligned}&(-1)^{2d-1}(-1)^{k+1}\beta\,dx_1\wedge\cdots\wedge dx_i\wedge dx_{k+2}'\wedge\cdots\wedge dx_d'\wedge dx_1'\\ &=-(-1)^{k+1}\beta\,dx_1\wedge\cdots\wedge dx_i\wedge dx_{k+2}'\wedge\cdots\wedge dx_d'\wedge dx_1'.\end{aligned}\tag{3.8}$$

Given these data, we compute $o^*(\overline{\mathscr{M}(f;r,q)})$ as follows.

$$\begin{aligned}o^*(\mathscr{M}_2(f;r,q))_x&=o^*(\mathscr{D}_r(f)\pitchfork\mathscr{A}_q(f))_x\\ &=o^*(\mathscr{D}_r(f))_x\wedge o^*(\mathscr{A}_q(f))_x\\ &=\alpha\beta\,dx_{k+2}\wedge\cdots\wedge dx_d.\end{aligned}$$

So we have

$$o^*(\overline{\mathscr{M}(f;r,q)})_x = \alpha\beta\, dx_{k+2}\wedge\cdots\wedge dx_d \wedge(-dx_1),$$

where $-dx_1$ corresponds to the dual of $\mathrm{grad}_x f$. Therefore,

$$o^*(\mathscr{A}_r(f))_x \wedge o^*(\overline{\mathscr{M}(f;r,q)})_{x'} = -\beta\, dx_1\wedge\cdots\wedge dx_i \wedge dx'_{k+2}\wedge\cdots\wedge dx'_d \wedge dx'_1.$$

This agrees with $(-1)^{k+1} = (-1)^{d(r)-d(q)}$ of the orientation of $\partial\overline{\mathscr{A}}_q(f)_1$ induced from $o^*(\mathscr{A}_q(f))_{(x,x')}$ given above in (3.8).

Lemma 3.16. *The normal orientation on* $\partial\overline{\mathscr{A}}_q(f)_1\cap(U_a\times\overline{L}_r)$ *induced from* $o^*(\mathscr{A}_q(f))_{(x,x')}$ *agrees with* $(-1)^{d(r)-d(q)}o^*(\mathscr{A}_r(f)\cap U_a)_x\wedge o^*(\overline{\mathscr{M}(f;r,q)})_{x'}$*.*

3.7. **Orientation of $\partial\overline{\mathscr{M}}_2(f)$.** To describe orientation of $\overline{\mathscr{M}}_2(f)$, we need to describe orientations of $\overline{\mathscr{M}}_2(f;U,U')$ for all charts $U,U'\subset M$. It suffices to describe orientations of the basic cases $\overline{\mathscr{M}}_2(f;U,\overline{L}_r)$, $U\cap\mathscr{A}_r(f)\neq\emptyset$, since the orientations of any pieces $\overline{\mathscr{M}}_2(f;U,U')$ are canonically induced from the basic ones.

The orientation of $\partial\overline{\mathscr{M}}_2(f;U,\overline{L}_r)$ is as follows. Let $\varphi''_0 : (\mathbb{R}^i\setminus\{0\})\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times(S^{i-1}_{\sqrt{\varepsilon}}\times\mathbb{R}^{d-i})$ and $\varphi'' : \widetilde{\gamma}^1(\mathbb{R}^i)\times\mathbb{R}^{d-i}\to\mathbb{R}^d\times(S^{i-1}_{\sqrt{\varepsilon}}\times\mathbb{R}^{d-i})$ be the same as above (in (3.5)). Then as above we have

$$\begin{aligned}\overline{\mathscr{M}}_2(f;U,\overline{L}_r) &= \mathrm{Im}\,\varphi''\cap(U\times\overline{L}_r)\text{ and}\\ \partial\,\mathrm{Im}\,\varphi'' &= (\{0\}\times\mathbb{R}^{d-i})\times(S^{i-1}_{\sqrt{\varepsilon}}\times\{0\}).\end{aligned}$$

Let $b=(\sqrt{\varepsilon},0,\ldots,0)\in\overline{L}_r$ and we consider the orientations on a neighborhood $\widetilde{U}_b$ of $(\mathscr{A}_r(f)\cap U)\times\{b\}\subset U\times\overline{L}_r$. We put

$$o(\mathscr{A}_r(f))_x = \alpha\, dx_{i+1}\wedge\cdots\wedge dx_d,$$

for $x\in M_r$, $\alpha,\beta\in\{+1,-1\}$, where these make sense. This gives

$$\begin{aligned}o^*(\mathscr{D}_r(f))_x &= \alpha\, dx_{i+1}\wedge\cdots\wedge dx_d,\\ o^*(\mathscr{A}_r(f))_x &= \alpha\, dx_1\wedge\cdots\wedge dx_i.\end{aligned}$$

We also consider $\overline{\mathscr{M}}_2(f;U,\overline{L}_r)$ as a submanifold of $\mathbb{R}^d\times\mathbb{R}^d$ via the embedding φ''_0. The orientation of $\overline{\mathscr{M}}_2(f;U,\overline{L}_r)$, considered as a submanifold of $\mathbb{R}^d\times\mathbb{R}^d$, induced from the standard orientation $dx_1\wedge\cdots\wedge dx_d$ of U is

$$o(\overline{\mathscr{M}}_2(f;U,\overline{L}_r))_{(x,x')} = dx'_1\wedge\cdots\wedge dx'_i\wedge dx_{i+1}\wedge\cdots\wedge dx_d.$$

Hence we obtain

$$o^*(\overline{\mathscr{M}}_2(f;U,\overline{L}_r))_{(x,x')} = -dx_1\wedge\cdots\wedge dx_i\wedge dx'_{i+1}\wedge\cdots\wedge dx'_d.$$

This induces the following orientation on the boundary:

$$\begin{aligned}&-(-1)^{2d-1}dx_1\wedge\cdots\wedge dx_i\wedge dx'_{i+1}\wedge\cdots\wedge dx'_d\wedge dx'_1\\ =&dx_1\wedge\cdots\wedge dx_i\wedge dx'_{i+1}\wedge\cdots\wedge dx'_d\wedge dx'_1\end{aligned}$$

for x' close to $b=(\sqrt{\varepsilon},0,\ldots,0)$.

Now the normal orientation of $\mathscr{D}_r(f)\cap\overline{L}_r$ is as follows.

$$\begin{aligned}o^*(\mathscr{D}_r(f)\cap\overline{L}_r)_{x'} &= o^*(\mathscr{D}_r(f))_{x'}\wedge(-dx'_1)\\ &= -\alpha\, dx'_{i+1}\wedge\cdots\wedge dx'_d\wedge dx'_1.\end{aligned}$$

Therefore,

$$o^*(\mathscr{A}_r(f))_x \wedge o^*(\mathscr{D}_r(f) \cap \overline{L}_r)_{x'} = -dx_1 \wedge \cdots \wedge dx_i \wedge dx'_{i+1} \wedge \cdots \wedge dx'_d \wedge dx'_1.$$

This agrees with -1 of the normal orientation of $\partial\overline{\mathscr{M}}_2(f;U,\overline{L}_r)$ induced from $o^*(\overline{\mathscr{M}}_2(f;U,\overline{L}_r))_{(x,x')}$ given above.

Lemma 3.17. *The normal orientation on $\partial\overline{\mathscr{M}}_2(f;U,\overline{L}_r)$ induced from $o^*(\overline{\mathscr{M}}_2(f;U,\overline{L}_r))_{(x,x')}$ agrees with $-o^*(\mathscr{A}_r(f) \cap U)_x \wedge o^*(\mathscr{D}_r(f) \cap \overline{L}_r)_{x'}$.*

3.8. **Orientation of $\overline{\mathscr{M}}_\Gamma$.**

3.8.1. *Standard orientation of $\mathscr{M}_\Gamma$ from graph orientation.* For a graph Γ without bivalent vertices, the moduli space $\mathscr{M}_\Gamma(\vec{f})$ can be defined as the intersection of submanifolds of $\mathrm{Conf}_n(M)$, as follows. Suppose for simplicity that the separated edges of Γ are labeled $1, 2, \ldots, a$. Let $\pi_{ij} : \mathrm{Conf}_n(M) \to \mathrm{Conf}_2(M)$ be the projection $(x_1, \ldots, x_n) \mapsto (x_i, x_j)$ and let Θ_k and H_k be the submanifolds of $\mathrm{Conf}_n(M)$ defined by

$$\Theta_k = \pi_{ij}^{-1}(\mathscr{M}_2(f_k)), \quad H_k = \pi_{ij}^{-1}(\mathscr{D}_{p_k}(f_k) \times \mathscr{A}_{q_k}(f_k)),$$

where $i = \sigma(k), j = \tau(k)$. Note that $\mathrm{codim}\,\Theta_k = d-1$, $\mathrm{codim}\,H_k = d-d(p_k)+d(q_k)$. Then we have

$$\mathscr{M}_\Gamma(\vec{f}) = \bigcap_{j=1}^{a} H_j \cap \bigcap_{j=a+1}^{m} \Theta_j,$$

where the intersection is transversal. Let $o^*(\Theta_j) \in \Omega_{\mathrm{dR}}^{d-1}(B)$ and $o^*(H_j) \in \Omega_{\mathrm{dR}}^{d-d(p_j)+d(q_j)}(B)$ be differential forms that represent normal orientations of Θ_j and H_j, and that are locally defined on a neighborhood B of a crossing point in $\mathrm{Conf}_n(M)$. We represent orientation of $\mathscr{M}_\Gamma(\vec{f})$ by a wedge product of $o^*(H_j)$'s and $o^*(\Theta_j)$'s.

Now we assume that $\Gamma \in \mathscr{G}_{n,m}(\vec{C})$, so that $\mathrm{codim}\,\Theta_j$ and $\mathrm{codim}\,H_j$ are even. We shall define a standard orientation of $\mathscr{M}_\Gamma(\vec{f})$ from the orientation of Γ, as follows. Recall that the orientation $o(\Gamma)$ of Γ is an orientation of the real vector space $\mathbb{R}^{B(\Gamma)} \oplus \bigoplus_{e \in \mathrm{Comp}(\Gamma)} \mathbb{R}^{H(e)}$ and we write $o(\Gamma) = o(\mathbb{R}^{B(\Gamma)}) \wedge \bigwedge_e o(\mathbb{R}^{H(e)})$. The $o(\mathbb{R}^{B(\Gamma)})$ part determines an orientation of $\mathrm{Conf}_n(M)$ and the $\bigwedge_e o(\mathbb{R}^{H(e)})$ part together with the orientations of the descending manifolds determine $o^*(\Theta_j)$'s and $o^*(H_j)$'s. Then we define the standard orientation of $\mathscr{M}_\Gamma(\vec{f})$ by the formula

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = \bigwedge_{j=1}^{a} o^*(H_j) \wedge \bigwedge_{j=a+1}^{m} o^*(\Theta_j) \in \Omega_{\mathrm{dR}}^{md-\sum_{j=1}^m \eta_j}(B).$$

Since $\mathrm{codim}\,\Theta_j$ and $\mathrm{codim}\,H_j$ are even, the order of wedge product does not matter.

The same rule equally works for graphs in $\mathscr{G}_{n,m,(2,1,\ldots,1)}(\vec{C})$, $\mathscr{G}_{n,m,(0,1,\ldots,1)}(\vec{C})$ etc. without bivalent vertices, since in that case only one H_j is odd codimensional, so again the orientation of $\mathscr{M}_\Gamma(\vec{f})$ is canonically determined from the graph orientation.

For a graph in $\mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ without bivalent vertices such that there is exactly one j with $\eta_j = 2$, exactly one j with $\eta_j = 0$ and otherwise $\eta_j = 1$, let j_2 and j_0 be such that $\eta_{j_2} = 2$, $\eta_{j_0} = 0$. Then we define a standard orientation of $\mathscr{M}_\Gamma(\vec{f})$ by

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = o^*(H_{j_0}) \wedge o^*(H_{j_2}) \wedge \bigwedge_{\substack{1\leq j\leq a\\ j\neq j_0, j_2}} o^*(H_j) \wedge \bigwedge_{j=a+1}^{m} o^*(\Theta_j).$$

We also consider orientations of $\mathscr{M}_\Gamma(\vec{f})$ for graphs $\Gamma \in \mathscr{G}_{n,m,(1,1,\ldots,1)}(\vec{C})$ with only one bivalent vertex. For three possiblilies for the position of the bivalent vertex in Γ, we define a standard orientation of $\mathscr{M}_\Gamma(\vec{f})$ as follows.

(1) $\Gamma =$ 

, $d(p_i) = d(r_i) + 1$. Let Γ' be the graph obtained from Γ by removing the segment $\partial^{(i)}(p_i, r_i)$. In this case, $\mathscr{M}_\Gamma(\vec{f})$ is canonically diffeomorphic to $\overline{\mathscr{M}(f_i; p_i, r_i)} \times \mathscr{M}_{\Gamma'}(\vec{f})$. We define

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = o^*(\overline{\mathscr{M}(f_i; p_i, r_i)}) \wedge o^*(\mathscr{M}_{\Gamma'}(\vec{f})).$$

(2) $\Gamma =$ 

, $d(q_i) = d(s_i) - 1$. Let Γ' be the graph obtained from Γ by removing the segment $\partial^{(i)}(s_i, q_i)$. In this case, $\mathscr{M}_\Gamma(\vec{f})$ is canonically diffeomorphic to $\mathscr{M}_{\Gamma'}(\vec{f}) \times \overline{\mathscr{M}(f_i; s_i, q_i)}$. We define

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = o^*(\mathscr{M}_{\Gamma'}(\vec{f})) \wedge o^*(\overline{\mathscr{M}(f_i; s_i, q_i)}).$$

(3) $\Gamma =$ 

. In this case, the orientation of $\mathscr{M}_\Gamma(\vec{f})$ is determined by the intersection of $H_k' = \pi_{ij}^{-1}(\mathscr{D}_{r_k}(f_k) \times \mathscr{A}_{r_k}(f_k))$ (codimension $(d - d(r_k)) + d(r_k) = d$) with the intersection of Θ_j's and H_j's. We define $o^*(\mathscr{M}_\Gamma(\vec{f}))$ as

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = o^*(H_{j_2}) \wedge \bigwedge_{\substack{1\leq j\leq a\\ j\neq j_2}} o^*(H_j) \wedge o^*(H_k') \wedge \bigwedge_{\substack{a+1\leq j\leq m\\ j\neq k}} o^*(\Theta_j). \tag{3.9}$$

3.8.2. *Induced orientation on $\partial\overline{\mathscr{M}}_\Gamma$.* Suppose that $\Gamma \in \mathscr{G}_{n,m,\Sigma(1,1,\ldots,1)}(\vec{C})$ has no bivalent vertices. We shall compare the orientation of a face of $\partial\overline{\mathscr{M}}_\Gamma(\vec{f})$ induced from $o^*(\mathscr{M}_\Gamma(\vec{f}))$ and the standard orientation of the same face of $\partial\overline{\mathscr{M}}_\Gamma(\vec{f})$ fixed in §3.8.1.

Suppose that $\Gamma \in \mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ has no bivalent vertex and that there is only one number j with $\eta_j = 2$ and $\eta_k = 1$ for $k \neq j$. Let j_2 be such that $\eta_{j_2} = 2$. As in §3.8.1, we consider $\mathscr{M}_\Gamma(\vec{f})$ as the intersection of the chains H_j's and Θ_j's in $\mathrm{Conf}_n(M)$.

Choose a number k such that $1 \leq k \leq m$ and put

$$\Sigma_k = \begin{cases} \bigcap_{\substack{1\leq j\leq a\\ j\neq k}} H_j \cap \bigcap_{j=a+1}^{m} \Theta_j & \text{if } 1 \leq k \leq a \\ \bigcap_{j=1}^{a} H_j \cap \bigcap_{\substack{a+1\leq j\leq m\\ j\neq k}} \Theta_j & \text{if } a+1 \leq k \leq m \end{cases}$$

Then

$$\begin{aligned}\operatorname{codim}\Sigma_k &= \operatorname{codim}\mathscr{M}_\Gamma(\vec{f}) - \operatorname{codim} H_k\\ &= (dn - d(n-m) - m - 1) - (d + \eta_k) = (d-1)(m+1) - 2d - \eta_k.\end{aligned}$$

Let X be one of the graphs that appear in the sum $d'_{\beta(k)}\Gamma$. We shall describe the orientation on the face $\mathscr{S}_X$ of $\partial\overline{\mathscr{M}}_\Gamma(\vec{f})$ corresponding to X, induced from the standard orientation of $\mathscr{M}_\Gamma(\vec{f})$.

(1) $X =$ 

. By Lemma 3.15, the orientation of $\mathscr{F}_{r_k}\overline{\mathscr{M}}_\Gamma(\vec{f})$ induced from the standard one

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = o^*(\mathscr{D}_{p_k}(f_k)) \wedge o^*(\mathscr{A}_{q_k}(f_k)) \wedge o^*(\Sigma_k)$$

is given by

$$\begin{aligned}&o^*(\overline{\mathscr{M}(f_k; p_k, r_k)}) \wedge o^*(\mathscr{D}_{r_k}(f_k)) \wedge o^*(\mathscr{A}_{q_k}(f_k)) \wedge o^*(\Sigma_k)\\ &= o^*(\overline{\mathscr{M}(f_k; p_k, r_k)}) \wedge o^*(\mathscr{M}_{\Gamma'}(\vec{f})).\end{aligned}$$

This agrees with the standard orientation fixed in §3.8.1.

(2) $X =$ 

. By Lemma 3.16, the induced orientation of $\mathscr{F}_{s_k}\overline{\mathscr{M}}_\Gamma(\vec{f})$ induced from the standard one

$$o^*(\mathscr{M}_\Gamma(\vec{f})) = o^*(\Sigma_k) \wedge o^*(\mathscr{D}_{p_k}(f_k)) \wedge o^*(\mathscr{A}_{q_k}(f_k))$$

is given by

$$\begin{aligned}&(-1)^{d(s_k)-d(q_k)} o^*(\Sigma_k) \wedge o^*(\mathscr{D}_{p_k}(f_k)) \wedge o^*(\mathscr{A}_{s_k}(f_k)) \wedge o^*(\overline{\mathscr{M}(f_k; s_k, q_k)})\\ &= (-1)^{d(s_k)-d(q_k)}(-1)^{(d-d(p_k)+d(q_k)+1)\operatorname{codim}\Sigma_k}\\ &\qquad \times o^*(\mathscr{D}_{p_k}(f_k)) \wedge o^*(\mathscr{A}_{s_k}(f_k)) \wedge o^*(\Sigma_k) \wedge o^*(\overline{\mathscr{M}(f_k; s_k, q_k)})\\ &= (-1)^{d(p_k)-d(s_k)} o^*(\mathscr{M}_{\Gamma'}(\vec{f})) \wedge o^*(\overline{\mathscr{M}(f_k; s_k, q_k)}).\end{aligned}$$

This differs from the standard orientation by $(-1)^{d(p_k)-d(s_k)}$.

(3) $X =$ 

. The induced orientation on the boundary is as in Lemma 3.17, which agrees with -1 of the standard orientation (3.9).

Now we have seen that the signs in the formula of the definition of d' is consistent with the induced orientations on the boundary of $\overline{\mathscr{M}}_\Gamma(\vec{f})$. Namely, we have the following proposition.

Proposition 3.18. *Suppose that $d = 3$. For $\Gamma \in \mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$ without bivalent vertices, we have*

$$\partial\overline{\mathscr{M}}_\Gamma(\vec{f}) = \overline{\mathscr{M}}_{(d+d')\Gamma}(\vec{f}) \coprod (\partial^{\mathrm{hi}}\overline{\mathrm{Conf}}_{2k}(M) \cap \overline{\mathscr{M}}_\Gamma(\vec{f}))$$

as oriented 0-dimensional manifolds.

3.9. **Moduli space of infinitesimal graph flows.** Fix a smooth Riemannian manifold X and a graph Γ with only compact edges that is a subgraph of a graph in $\mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$. We consider the moduli space of a singular graph with Γ collapsed into a point. Since Γ is a subgraph of a graph in $\mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$, it may have vertices of valence ≤ 2 that is a part of a black vertex. We call such a vertex a bivalent or a univalent black vertex. Suppose that Γ has n vertices and m edges. We consider the trivial bundles

$$\begin{aligned} \pi &: X \times S^{d-1} \to X, \\ \pi' &: X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d) \to X. \end{aligned}$$

Note that X is not necessarily d dimensional. Let $\phi_{ij} : \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d) \to S^{d-1}$ be the Gauss map that assigns the unit vector $\frac{x_j - x_i}{\|x_j - x_i\|}$ and let

$$\widetilde{\phi}_{ij} : X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d) \to X \times S^{d-1}$$

be the map defined by $\widetilde{\phi}_{ij}(y, x) = (\phi_{ij}(y), x)$. Suppose given a smooth section $\gamma : X \to X \times S^{d-1}$ of π. Since $\widetilde{\phi}_{ij}$ is obviously transversal to $\gamma(X)$ on each fiber, the subset

$$\Theta_k(\gamma) = \widetilde{\phi}_{ij}^{-1}(\gamma(X)) \subset X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d)$$

forms a smooth subbundle of π'. Now we define

$$\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma}) = \bigcap_{k=1}^{m} \Theta_k(\gamma_k) \subset X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d) \tag{3.10}$$

for a sequence $\vec{\gamma} = (\gamma_1, \ldots, \gamma_m)$ of sections of the sphere bundle π. One can also define $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ for a section $\vec{\gamma}$ of the trivial $\mathbb{R}^d - \{0\}$-bundle over X similarly. The formula (3.10) also defines an orientation of $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ if the intersection is transversal.

Lemma 3.19. *For a generic choice of $\vec{\gamma}$, the moduli space $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ is a smooth submanifold of $X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d)$ of codimension $(d-1)m$. If X is compact, then so is $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$.*

Proof. Note that $\Theta_k(\gamma)$ is a smooth submanifold of codimension $d-1$. By the transversality theorem, one can deform $\vec{\gamma}$ slightly and inductively so that the intersection (3.10) is transversal in the above sense. Thus for a generic choice of $\vec{\gamma}$, $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ is a smooth submanifold. □

Lemma 3.20. *Suppose that Γ has a bivalent black vertex a and that the edges of Γ including the vertex a are (b, a) and (a, c). Let Γ' be the labeled graph obtained from Γ by exchenging labels for edges (b, a) and (a, c). Then there is an involution $S : X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d) \to X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d)$ which leaves $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma}) \coprod \mathscr{M}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma})$ invariant and reverses the orientation.*

Proof. Let $(t, x_1, \ldots, x_n) \in X \times \overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d)$ be a point on $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$. Suppose that the vertices a, b and c of Γ correspond to x_α, x_β and x_γ respectively. Then the automorphism of $\overline{\mathrm{Conf}}_n^{\mathrm{local}}(\mathbb{R}^d)$ which sends x_α to $x_\beta + x_\gamma - x_\alpha$ gives rise to the desired involution. □

Lemma 3.21. *Suppose that Γ has a univalent black vertex a and $|V(\Gamma)| \geq 3$. Then $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ admits a smooth free $\mathbb{R}$-action.*

Proof. Suppose that the edge of Γ including the vertex a is (a,b). Since $|V(\Gamma)| \geq 3$, there is a dilation of the trajectory corresponding to (a,b) in a graph flow, which is independent of overall dilation. This gives a desired $\mathbb{R}$-action. $\square$

Now we consider the case $d = 3$, $\Gamma \in \mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$, in which case $\overline{\mathscr{M}}_\Gamma(\vec{f})$ is 1-dimensional by Proposition 3.4. For a subset A of $\{1,\ldots,n\}$, let Γ_A denote the subgraph of Γ with $V(\Gamma_A) = \alpha(A)$ such that $E(\Gamma_A) = \mathrm{Comp}(\Gamma_A)$. Let $\vec{f}_A \subset \vec{f}$ be the subsequence corresponding to $E(\Gamma_A)$. According to the description of $\partial\overline{\mathrm{Conf}}_n(M)$ in §3.4, the face of $\partial\overline{\mathscr{M}}_\Gamma(\vec{f})$ corresponding to the face $\partial_A\overline{\mathrm{Conf}}_n(M)$ is diffeomorphic via the induced map by a trivialization $TM \overset{\approx}{\to} M \times \mathbb{R}^d$ (framing), to

$$\overline{\mathscr{M}}_{\Gamma/\Gamma_A}(\vec{f} \setminus \vec{f}_A) \times \mathscr{M}_{\Gamma_A}^{\mathrm{local}}(-\mathrm{grad}_0 j^1 \vec{f}_A),$$

which is at most a 0-dimensional oriented manifold. By Lemma 3.20, we have $\#\mathscr{M}_{\Gamma_A}^{\mathrm{local}}(-\mathrm{grad}_0 j^1 \vec{f}_A) + \#\mathscr{M}_{\Gamma'_A}^{\mathrm{local}}(-\mathrm{grad}_0 j^1 \vec{f}_A) = 0$ if Γ_A has a bivalent black vertex. By Lemma 3.21, if $|A| \geq 3$ and if Γ_A has a univalent vertex, then

$$\mathscr{M}_{\Gamma_A}^{\mathrm{local}}(-\mathrm{grad}_0 j^1 \vec{f}_A) \cong \emptyset \times \mathbb{R} = \emptyset.$$

Since $\Gamma \in \mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$, the subgraph Γ_A must have bivalent or univalent black vertices. Thus Proposition 3.18 implies the following lemma.

Proposition 3.22. *Suppose that $d = 3$, $\Gamma \in \mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$. Then we have*

$$\#\overline{\mathscr{M}}_{(d+d')\Gamma}(\vec{f}) = \#\partial\overline{\mathscr{M}}_\Gamma(\vec{f}) = 0.$$

3.10. **Definition of $\widehat{Z}_{2k,3k}$.** Now we assume that $d = 3$, $n = 2k$, $m = 3k$.

3.10.1. *Definition of $Z_{2k,3k}$.* By Proposition 3.4, for any graph $\Gamma \in \mathscr{G}_{2k,3k}(\vec{C})$ without bivalent vertices the moduli space $\overline{\mathscr{M}}_\Gamma(\vec{f})$ is a compact 0-dimensional submanifold of $\overline{\mathrm{Conf}}_{2k}(M)$. So the number of the components can be counted with signs if we orient $\overline{\mathscr{M}}_\Gamma(\vec{f})$ by the standard orientation of §3.8. Let $\#\overline{\mathscr{M}}_\Gamma(\vec{f})$ be the number of components counted with signs. Then the correspondence $\Gamma \mapsto \#\overline{\mathscr{M}}_\Gamma(\vec{f})$ defines a linear map

$$\#\overline{\mathscr{M}}_\bullet(\vec{f}) : \mathscr{G}_{2k,3k}(\vec{C}) \to \mathbb{Q}.$$

Lemma 3.23. *Let $\mathscr{H}'_{2k}(\vec{C})$ be the image of $\mathscr{G}_{2k,3k}(\vec{C})$ under $\langle\cdot\rangle : \mathscr{G}_{2k,3k}(\vec{C}) \to \mathscr{H}_{2k}(\vec{C})$. There is a unique linear map*

$$\#\overline{\mathscr{M}}'_\bullet(\vec{f}) : \mathscr{H}'_{2k}(\vec{C}) \to \mathbb{Q}$$

such that $\#\overline{\mathscr{M}}_\bullet(\vec{f}) = \#\overline{\mathscr{M}}'_\bullet(\vec{f}) \circ \langle\cdot\rangle$.

Proof. That the ∂-relation and C-relation are included in the kernel of $\#\overline{\mathscr{M}}_\bullet(\vec{f})$ is obvious from the definition of the moduli space $\overline{\mathscr{M}}_\Gamma(\vec{f})$ and the Morse complex. We show that $(d+d')\mathscr{G}_{2k,3k,\Sigma(1,\ldots,1)}(\vec{C})$ is included in the kernel of $\#\overline{\mathscr{M}}_\bullet(\vec{f})$. But this follows immediately from Proposition 3.22. The proof is completed. $\square$

Now we define

$$\begin{aligned} Z_{2k,3k}(\vec{f}) &= \#\overline{\mathscr{M}}'_{\bullet}(\vec{f})\,\langle\widetilde{\mathrm{Tr}}'_{\vec{g}}(\widetilde{\gamma}_{2k,3k})\rangle \\ &= \sum_{\Gamma\in\mathscr{G}_{2k,3k}(\vec{C})} \widetilde{\mathrm{Tr}}_{\vec{g}}[[\Gamma]]\cdot\#\overline{\mathscr{M}}_{\Gamma}(\vec{f})\in\mathscr{A}_{2k,3k}. \end{aligned}$$

By Lemma 2.3, $Z_{2k,3k}(\vec{f})$ does not depend on the choice of $\vec{g}$. However it may still depend on the choice of $\vec{f}$. We need to consider a correction term to make this an invariant of M.

3.10.2. *Differential relation $\mathscr{H}$.* Let $\mathscr{H}\subset J^3(\mathbb{R}^d,\mathbb{R})$ be the subspace of all $f\in J^3(\mathbb{R}^d,\mathbb{R})$ such that one of the following holds:

1. $0\in\mathbb{R}^d$ is not a critical point of f,
2. $0\in\mathbb{R}^d$ is a nondegenerate critical point of f ($\Leftrightarrow$ $0\in\Sigma^1(f)$),
3. $0\in\mathbb{R}^d$ is a birth/death singularity of f ($\Leftrightarrow$ $0\in\Sigma^2(f)$).

Since $\mathscr{H}$ is $GL_+(\mathbb{R}^d)$-invariant, we can consider the associated subbundle $\mathscr{H}(TX)$ of the jet bundle $J^3(X,\mathbb{R})$ with the structure group $GL_+(\mathbb{R}^d)$ if X is an oriented smooth d-manifold. By K. Igusa's lemma saying that the "framed" version of $\mathscr{H}$ is weakly contractible ([Ig1]), one obtains the following lemma immediately.

Lemma 3.24. *Let M be a closed oriented d-manifold and let W be a compact oriented $(d+1)$-manifold with $\partial W=M$. Suppose that there exists a rank d subbundle T^vW of TW with $T^vW|_M=TM$. Then any section $\gamma\in\Gamma(\mathscr{H}(TM))$ has an extension $\widetilde{\gamma}\in\Gamma(\mathscr{H}(T^vW))$.*

3.10.3. *Correction term.* Now we shall define a correction term. We consider S^3 as the one point compactification $\mathbb{R}^3\cup\{\infty\}$. Let

$$U_\infty=\{x\in\mathbb{R}^3\,;\,\|x\|>R\}\cup\{\infty\}\subset S^3$$

for some large R. Choose a compact spin 4-manifold W with $\partial W=M$ and choose a point $\infty\in M$ and a small open ball $U'_\infty\subset M$ including ∞. Fix a diffeomorphism $\varphi_\infty:U'_\infty\to U_\infty$ which sends ∞ to ∞. Since $\Omega_3^{\mathrm{spin}}=0$, if a framing τ_M on $T(M\setminus\{\infty\})$ is given, then one can find a compact spin 4-manifold W with $\partial W=M$ and with a spin structure that is compatible with the spin structure of M determined by τ_M. We fix τ_M such that it agrees on $U'_\infty\setminus\{\infty\}$ with the pullback of the standard one on $U_\infty\setminus\{\infty\}$ by φ_∞. We extend the trivial stabilization $n\oplus\tau_M$ on $\varepsilon^1\oplus T(M\setminus\{\infty\})$ to a 4-framing τ'_M on $\varepsilon^1\oplus TM$, which is possible since S^3 is stably parallelizable.

Lemma 3.25 (Lemma 2.3 of [KM]). *Suppose that τ'_M is compatible with the spin structure on W. Then τ'_M extends to a framing on W if and only if $p_1(TW;\tau'_M)=0$ and $\chi(W)=1$, where $p_1(TW;\tau'_M)\in\mathbb{Z}$ denotes the relative Pontrjagin number.*

Lemma 3.26 (Lemma 2.3 of [KM]). *We have $p_1(TW;\tau'_M)\equiv 0$ mod 2. Moreover, $p_1(TW;\tau'_M)$ can be changed by ± 2 by changing τ_M by a fiberwise SO_3-action given by a section $M\to SO_3$.*

By Lemma 3.26, there is a unique framing τ_M with $p_1(TW;\tau'_M)=0$. Moreover, since T^4 is a parallelizable spin 4-manifold and since $\chi(W\#T^4)=\chi(W)-2$, one can find a pair (W, τ'_M) satisfying the condition of Lemma 3.25, if $\chi(W) \equiv 1 \bmod 2$ and $\chi(W) > 0$. The following lemma follows from the proof of [KM, Theorem 2.6].

Lemma 3.27. *For a $\mathbb{Z}$-homology 3-sphere M (with a canonical spin structure), there exists a simply-connected compact spin 4-manifold W with $\partial W = M$ as the spin boundary such that $\chi(W) \equiv 1$ mod 2 and $\chi(W) > 0$.*

There is another extension of τ_M that is canonical. The standard framing $\tau_{\mathbb{R}^3}$ on $\mathbb{R}^3$ can be replaced within $U_\infty \setminus \{\infty\}$ so that it extends to a framing of TS^3. We denote the resulting framing of TS^3 by τ_{S^3}. Moreover τ_{S^3} can be choosen so that the signature defect vanishes:

$$\frac{1}{3}p_1(TD^4; \tau_{S^3} \oplus n) - \operatorname{sign} D^4 = 0.$$

Such a choice is canonical up to homotopy ([KM]). Then by gluing the pullback $\varphi_\infty^*(\tau_{S^3}|_{U_\infty})$ with $\tau_M|_{M\setminus U'_\infty}$, we obtain a honest framing $\overline{\tau}_M$ of TM. The two framings $\overline{\tau}_M$ and τ'_M define the bundle monomorphism

$$\theta_M = \tau_M'^{-1} \circ \overline{\tau}_M : TM \to TW|_M = \varepsilon^1 \oplus TM.$$

Note that θ_M is unique up to homotopy in the sense that if one changes the honest framing τ_M within $T(M \setminus U'_\infty)$ by a map $(M \setminus U'_\infty, \partial(M \setminus U'_\infty)) \to (GL(\mathbb{R}^3), \mathrm{id})$, then $\overline{\tau}_M$ and τ'_M may change while θ_M will not change.

Now let (W, τ'_M) be a pair satisfying the condition of Lemma 3.25. Then one can find a sub 3-framing τ_W^v of TW that extends τ_M and that is a sub 3-framing of τ'_M. The 3-framing τ_W^v spans a rank 3 subbundle T^vW of TW. Let $\mathscr{H}(T^vW)$ be the associated $\mathscr{H}$-bundle to T^vW. Then θ_M induces a bundle isomorphism $\theta_M : \mathscr{H}(TM) \to \mathscr{H}(T^vW)$. For each i, let ρ_i be a section of $\mathscr{H}(T^vW)$ given by $\theta_M(j^3 f_i)$ and put $\vec{\rho}_W = (\rho_1, \ldots, \rho_{3k})$. We define

$$Z^{\mathrm{anomaly}}_{2k,3k}(\vec{\rho}_W) = \sum_{\Gamma \in \mathscr{G}_{2k,3k}} [\Gamma] \cdot \#\mathscr{M}^{\mathrm{local}}_\Gamma(-\mathrm{grad}_0 j^1 \vec{\rho}_W) \in \mathscr{A}_{2k,3k},$$

where the moduli space $\mathscr{M}^{\mathrm{local}}_\Gamma(-\mathrm{grad}_0 j^1 \vec{\rho}_W)$ is considered inside the trivial $\overline{\mathrm{Conf}}^{\mathrm{local}}_{2k}(\mathbb{R}^3)$-bundle over $\bigcap_{j=1}^{3k}(W \setminus \Sigma(\rho_j))$ associated to the restriction of the $\mathbb{R}^3$-bundle T^vW.

Lemma 3.28. *For a generic choice of the extension $\vec{\rho}_W$, the number $\#\mathscr{M}^{\mathrm{local}}_\Gamma(-\mathrm{grad}_0 j^1 \vec{\rho}_W)$ is finite.*

Proof. We put $\vec{\gamma} = (\gamma_1, \ldots, \gamma_{3k}) = -\mathrm{grad}_0 j^1 \vec{\rho}_W$ for simplicity. Since $\Sigma(\gamma_i) \subset W$ is a smooth 1-dimensional submanifold of W, we may assume without loss of generality that $\Sigma(\gamma_i) \cap \Sigma(\gamma_j) = \emptyset$ if $i \neq j$. We shall show that the projection of $\mathscr{M}^{\mathrm{local}}_\Gamma(\vec{\gamma})$ on W is disjoint from a neighborhood of $\Sigma(\gamma_j)$ and hence from a neighborhood of $\coprod_{j=1}^{3k} \Sigma(\gamma_j)$. Let Γ' be the graph obtained from Γ by replacing $E(\Gamma)$ with $E(\Gamma) \setminus \{\beta(j)\}$. By a similar argument as in the proof of Lemma 3.19, one sees that $\mathscr{M}^{\mathrm{local}}_{\Gamma'}(\vec{\gamma} \setminus \{\gamma_j\})$ is a smooth submanifold of $W \times \overline{\mathrm{Conf}}^{\mathrm{local}}_{2k}(\mathbb{R}^3)$ of codimension $2(3k-1) = 6k-2$, i.e., a 2-dimensional submanifold. So, for a generic choice of $\vec{\gamma}$, the projection of $\mathscr{M}^{\mathrm{local}}_{\Gamma'}(\vec{\gamma} \setminus \{\gamma_j\})$ on W is disjoint from a neighborhood of $\Sigma(\gamma_j)$ due to a dimensional reason. Recall from definition that $\mathscr{M}^{\mathrm{local}}_\Gamma(\vec{\gamma})$ is the intersection of $\mathscr{M}^{\mathrm{local}}_{\Gamma'}(\vec{\gamma} \setminus \{\gamma_j\})$ with $\Theta_j(\gamma_j)$. Hence for a generic choice of $\vec{\gamma}$, the projection of $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ on W is disjoint from a neighborhood of $\Sigma(\gamma_j)$. By Lemma 3.19, the restriction of $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ to the complement of a neighborhood of $\coprod_{j=1}^{3k}\Sigma(\gamma_j)$ is compact. So, for a generic choice of $\vec{\gamma}$, $\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})$ is a compact 0-dimensional submanifold. □

We say that two compact spin 4-manifolds W and W' with $\partial W = \partial W' = M$ are *relatively spin cobordant* if there is a compact spin 5-manifold V with $\partial V = W \cup_\partial (-W')$ whose spin structure is an extension of those of W and $-W'$.

Proposition 3.29. *Let W and W' be two compact spin 4-manifolds with $\partial W = \partial W' = M$ and $\chi(W) = \chi(W') = 1$. If W and W' are relatively spin cobordant, then the following assertions hold.*

1. *There exists a framing τ_M of $T(M \setminus \{\infty\})$ such that τ'_M can be extended to 4-framings of both TW and TW'. Hence one can find sequences of sections $\vec{\rho}_W \in \Gamma(\mathscr{H}(T^vW))^m$ and $\vec{\rho}_{W'} \in \Gamma(\mathscr{H}(T^vW'))^m$ with $\vec{\rho}_W|_M = \vec{\rho}_{W'}|_M = \theta_M(j^3\vec{f})$.*
2. *For any such extensions $\vec{\rho}_W$ and $\vec{\rho}_{W'}$, which are generic in the sense of Lemma 3.28, we have*

$$Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W) = Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_{W'}). \tag{3.11}$$

Proof. (1) Put

$$X = W \cup_g (M \times [0,1]) \cup_{g'} (-W'),$$

where the gluing maps $g : \partial W = M \to M \times \{1\}$ and $g' : -\partial W = -M \to M \times \{0\}$ are the natural ones. By assumption, we have $[X] = 0 \in \Omega_4^{\mathrm{spin}}$. Then by the isomorphism $\mathrm{sign} : \Omega_4^{\mathrm{spin}} \xrightarrow{\sim} 16\mathbb{Z}$ and by Hirzebruch's signature theorem $\mathrm{sign}\, X = \frac{1}{3}\langle p_1(TX), [X]\rangle$ for X closed, it holds that

$$p_1(TW; \tau'_M) - p_1(TW'; \tau'_M) = 0$$

for any choice of τ_M. Hence by choosing τ_M suitably, we may assume that $p_1(TW; \tau'_M) = p_1(TW'; \tau'_M) = 0$ by Lemma 3.26. Thus by Lemma 3.25, such a τ'_M extends to 4-framings on both W and W'.

(2) One obtains a 4-framing τ_X on TX by gluing the framings on TW, TW' and $\mathrm{pr}_1^*\tau'_M$ on $T(M \times [0,1])$ that are compatible at $M \times \{0,1\}$. The sub 3-framing of τ_X that extends τ_M spans a rank 3 subbundle T^vX of TX. One obtains a section $\vec{\rho}_X$ of $\mathscr{H}(T^vX)$ by gluing $\vec{\rho}_W$ and $\vec{\rho}_{W'}$ together at the boundary. By definition of $Z_{2k,3k}^{\mathrm{anomaly}}$, we have

$$Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X) = Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W) - Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_{W'}). \tag{3.12}$$

Since X is spin null-cobordant, there exists a compact spin 5-manifold V with $\partial V = X$. We want a parallelizable one for such a V. We consider the obstruction to extending the stable framing $n \oplus \tau_X$ of $\varepsilon^1 \oplus TX$ over a 5-framing on TV. Since V is spin and since $\pi_2(SO_5) = 0$, $\pi_3(SO_5) = \mathbb{Z}$ and $\pi_4(SO_5) = \mathbb{Z}_2$, the first obstruction $\mathfrak{o}_1(V; n \oplus \tau_X)$ to the extension lies in the group $H^4(V, \partial V; \pi_3(SO_5)) \cong H_1(V; \mathbb{Z})$. It is easy to see that any class in $H_1(V; \mathbb{Z})$ can be realized by an embedding $f : S^1 \to \mathrm{Int}\, V$. Since V is spin, the normal bundle N_f of the image of f is trivial. So by a surgery along a framed embedding (f, τ_f), the homology class $[f]$ can be eliminated. Moreover, by choosing τ_f suitably, we may assume that the resulting 5-manifold of the surgery is spin again since $\pi_1 SO_4 \to \pi_1 SO_5$ is surjective.

Now we assume $H_1(V;\mathbb{Z})=0$ by doing surgeries as in the previous paragraph, if necessary. Then the next obstruction $\mathfrak{o}_2(V;n\oplus\tau_X)$ for the extension lies in the group $H^5(V,\partial V;\pi_4(SO_5))\cong\mathbb{Z}_2$. To eliminate $\mathfrak{o}_2(V;n\oplus\tau_X)$, we consider the connected sum $V'=V\#V$ between the interiors. Then one can check that the obstruction $\mathfrak{o}_2(V';n\oplus\tau_X\coprod n\oplus\tau_X)\in H^5(V',\partial V';\pi_4(SO_5))$ vanishes in any case. This change makes both sides of (3.11) just multiplied by 2. So we may assume without trouble that the obstruction $\mathfrak{o}_2(V;n\oplus\tau_X)$ vanishes in advance. Thus we assume that we have a framed 5-manifold (V,τ_V), that extends $(X,n\oplus\tau_X)$. Then a sub 3-framing of τ_V defines a rank 3 subbundle T^vV of TV, that is an extension of T^vX. Moreover there is an extension $\vec{\rho}_V=(\rho_{V,1},\ldots,\rho_{V,3k})\in\Gamma(\mathscr{H}(T^vV))$ of $\vec{\rho}_X$.

We put $\vec{\gamma}=(\gamma_1,\ldots,\gamma_{3k})=-\mathrm{grad}_0 j^1\vec{\rho}_V$. Let Γ' be the graph obtained from Γ by replacing $E(\Gamma)$ with $E(\Gamma)\setminus\{\beta(j)\}$. We identify the total space of the associated $\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$-bundle to T^vV with $V\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ via τ_V. Let $\pi' : V\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)\to V$ be the projection. Then as mentioned in the proof of Lemma 3.28, $\mathscr{M}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\})$ is a smooth submanifold of $V\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ of codimension $6k-2$, i.e., 3-dimensional. On the other hand, $\pi'^{-1}(\Sigma(\gamma_j))$ is a codimension 3 submanifold of $V\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$. For a generic choice of γ_j, the intersection of the two submanifolds is transversal and hence the intersection is a 0-dimensional submanifold.

Take a point $x\in\mathscr{M}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\})\pitchfork\pi'^{-1}(\Sigma(\gamma_j))$ and a small neighborhood U_x' of x in $V\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ so that U_x' contains exactly one intersection point. Let $U_x=\pi'(U_x')$. By a suitable small deformation of $\vec{\gamma}\setminus\{\gamma_j\}$, we may assume that

1. $\pi'(x)$ is a Morse point of $\rho_{V,j}$ and $U_x\cap\Sigma^2(\rho_{V,j})=\emptyset$,
2. $\pi'(\mathscr{M}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\}))$ is tangent to T^vV. (This is possible since $\Sigma(\gamma_j)$ is transversal to T^vV.)

We consider the blow-up $B\ell(U_x,U_x\cap\Sigma(\gamma_j))$ and consider the closure of

$$\mathscr{M}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\})\cap(\pi'^{-1}(U_x\setminus\Sigma(\gamma_j)))$$

in $B\ell(U_x,U_x\cap\Sigma(\gamma_j))\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$. We denote the closure by $\overline{\mathscr{M}}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\};U_x)$. Since $\mathscr{M}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\})$ is transversal to $\pi'^{-1}(\Sigma(\gamma_j)\cap U_x)$, the closure $\overline{\mathscr{M}}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\};U_x)$ is a smooth submanifold of $B\ell(U_x,U_x\cap\Sigma(\gamma_j))\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ with boundary. On the other hand, $\Theta_j(\gamma_j)\subset(V-\coprod_{\ell=1}^{3k}\Sigma(\gamma_\ell))\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ has the closure $\overline{\Theta}_j(\gamma_j;U_x)$ in $B\ell(U_x,U_x\cap\Sigma(\gamma_j))\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$, that is a smooth submanifold with boundary. Since $U_x\cap\Sigma(\gamma_j)$ consists only of Morse points, we may assume that the intersection of $\overline{\Theta}_j(\gamma_j;U_x)$ and $\overline{\mathscr{M}}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\};U_x)$ is transversal (even on the boundary) and forms a 1-dimensional smooth submanifold of $B\ell(U_x,U_x\cap\Sigma(\gamma_j))\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ with boundary. Now we define

$$\overline{\mathscr{M}}_{\Gamma}^{\mathrm{local}}(\vec{\gamma};U_x)=\overline{\Theta}_j(\gamma_j;U_x)\pitchfork\overline{\mathscr{M}}_{\Gamma'}^{\mathrm{local}}(\vec{\gamma}\setminus\{\gamma_j\};U_x).$$



Figure 4.

By the assumption (2), the boundary of $\pi'(\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x))$ lies in the fiber S_x^2 of the normal sphere bundle $S(T^vV)$ at $\pi'(x)$. Let Γ^* denote the graph obtained from Γ by reversing the orientation of the edge labeled j. Since $\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)\coprod\overline{\mathscr{M}}_{\Gamma^*}^{\mathrm{local}}(\vec{\gamma};U_x)$ is transversal to $\partial B\ell(U_x,U_x\cap\Sigma(\gamma_j))\times\overline{\mathrm{Conf}}_{2k}^{\mathrm{local}}(\mathbb{R}^3)$ and since on a neighborhood of $\Sigma(\gamma_j)$ there is a symmetry between the moduli spaces $\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)$ and $\overline{\mathscr{M}}_{\Gamma^*}^{\mathrm{local}}(\vec{\gamma};U_x)$ due to the symmetry of the standard model of a Morse point, the intersection of $\pi'(\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)\coprod\overline{\mathscr{M}}_{\Gamma^*}^{\mathrm{local}}(\vec{\gamma};U_x))$ with $\partial B\ell(U_x,U_x\cap\Sigma(\gamma_j))$ consists of two points in S_x^2 that are in an antipodal position by the assumption (1). Hence one can check that

$$\begin{aligned}
&[\Gamma]\cdot\#\partial\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)+[\Gamma^*]\cdot\#\partial\overline{\mathscr{M}}_{\Gamma^*}^{\mathrm{local}}(\vec{\gamma};U_x)\\
=&[\Gamma]\Big(\#\partial\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)-\#\partial\overline{\mathscr{M}}_{\Gamma^*}^{\mathrm{local}}(\vec{\gamma};U_x)\Big)\\
=&[\Gamma]\Big(\#\partial\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)-\#\partial\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)\Big)=0
\end{aligned}$$

since the symmetry reverses the orientation of $\overline{\Theta}_j$, and the inward normal vectors at $\#\partial\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(\vec{\gamma};U_x)$ and $\#\partial\overline{\mathscr{M}}_{\Gamma^*}^{\mathrm{local}}(\vec{\gamma};U_x)$ are opposite. See Figure 4. Hence we have

$$\begin{aligned}
0&=\sum_{\Gamma\in\mathscr{G}_{2k,3k}}[\Gamma]\cdot\#\partial\mathscr{M}_\Gamma^{\mathrm{local}}(\vec{\gamma})\\
&=Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X)-\sum_{\Gamma\in\mathscr{G}_{2k,3k}}[\Gamma]\cdot\#\mathscr{M}_{d\Gamma}^{\mathrm{local}}(\vec{\gamma}).
\end{aligned}$$

By the relation of $\mathscr{A}_{2k,3k}$, the second term in the second row vanishes. This together with (3.12) gives the desired identity. □

Next we consider general pairs of spin 4-manifolds W, W' with $\chi(W)=\chi(W')=1$ which may not be relatively spin cobordant. We choose 3-framings σ_M and τ_M on $T(M\setminus\{\infty\})$ so that

$$p_1(TW;\tau_M')=p_1(TW';\sigma_M')=0.$$

Then by Lemma 3.25, τ_M' extends to a 4-framing of W and σ_M' extends to a 4-framing of W'. But in general τ_M may not be homotopic to σ_M, so we may not have a 4-framing of $X=W\cup_g(M\times[0,1])\cup_{g'}(-W')$, namely, X may be just almost parallelizable. Although we do not have a framing of X, we have a natural rank 3 (possibly nontrivial) subbundle T^vX of TX that agrees with $\mathrm{pr}_1^*T(M\setminus\{\infty\})$ on $(M\setminus\{\infty\})\times[0,1]$, which extends those determined by σ_M and τ_M. By choosing a generic section $\vec{\rho}_X\in\Gamma(\mathscr{H}(T^vX))$, one can also define $Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X)\in\mathscr{A}_{2k,3k}$. By a similar argument, one can define $Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X)$ for any almost parallelizable, closed, connected, spin 4-manifold with $\chi(X)=2$. Let $\Omega_4^{\mathrm{spin}}(2)$ denote the set of spin cobordism classes of almost parallelizable, closed, connected, spin 4-manifolds X with $\chi(X)=2$. By Proposition 3.29, the correspondence $X\mapsto Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X)$ for generic $\vec{\rho}_X$ defines a well-defined map

$$Z_{2k,3k}^{\mathrm{anomaly}}:\Omega_4^{\mathrm{spin}}(2)\to\mathscr{A}_{2k,3k}.$$

The set $\Omega_4^{\mathrm{spin}}(2)$ has a group structure given by connected sum. More precisely, if X is an almost parallelizable, closed, connected, spin 4-manifold with $\chi(X)=2$, then there is a framing on $X-D^4$. If X' is another almost parallelizable, closed, connected, spin 4-manifold with $\chi(X')=2$, then by forming the boundary connected sum $\overline{X-D^4}\natural\overline{X'-D^4}$ and capping by D^4 along the boundary in a natural way, we will obtain an almost parallelizable, closed, conneced, spin 4-manifold X'' with $\chi(X'')=2$ that is diffeomorphic to $X\#X'$. This defines an abelian group structure on $\Omega_4^{\mathrm{spin}}(2)$ on which the inverse of X is given by $-X$.

Lemma 3.30. *The map $Z_{2k,3k}^{\mathrm{anomaly}}:\Omega_4^{\mathrm{spin}}(2)\to\mathscr{A}_{2k,3k}$ is a group homomorphism.*

Proof. If $[X]=0\in\Omega_4^{\mathrm{spin}}(2)$, then we have $0=\operatorname{sign}X=\frac{1}{3}\langle p_1(TX),[X]\rangle$ so the stabilization of the 4-framing on $\partial\overline{X-D^4}$ induced from that of $X-D^4$ extends over D^4. This implies that X is stably parallelizable. Then the proof of Proposition 3.29 shows that $Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X)=0$ for any generic $\vec{\rho}_X\in(\Gamma(\mathscr{H}(T^vX)))^{3k}$. The additivity of $Z_{2k,3k}^{\mathrm{anomaly}}$ follows from the fact that $Z_{2k,3k}^{\mathrm{anomaly}}$ is spin cobordism invariant, as shown in the proof of Proposition 3.29, and that $X\coprod X'$ and $X\#X'$ are spin cobordant. Hence $Z_{2k,3k}^{\mathrm{anomaly}}$ is a homomorhism. □

Proposition 3.31. *Let W and W' be compact, connected, spin 4-manifolds with $\partial W=\partial W'=M$, $\chi(W)=\chi(W')=1$ and suppose that $\vec{\rho}_W|_M=\vec{\rho}_{W'}|_M$. Then for each $k\geq 1$ there exists a constant $\mu_k\in\mathscr{A}_{2k,3k}$ such that*

$$Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W)-\mu_k\operatorname{sign}W=Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_{W'})-\mu_k\operatorname{sign}W'.$$

Hence $Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W)-\mu_k\operatorname{sign}W$ does not depend on the choice of $(W,\vec{\rho}_W)$ such that $\partial W=M$, $\chi(W)=1$, $\vec{\rho}_W|_{\partial M}=\theta_M(j^3\vec{f})$.

Proof. By Lemma 3.30, $Z_{2k,3k}^{\mathrm{anomaly}}$ is a restriction of a homomorphism $\Omega_4^{\mathrm{spin}}\to\mathscr{A}_{2k,3k}$. So there exists a constant $\mu_k\in\mathscr{A}_{2k,3k}$ such that

$$Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_X)=\mu_k\operatorname{sign}X,$$

for $X=W\cup_g(M\times[0,1])\cup_{g'}(-W')$. Hence

$$\begin{aligned}Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W)-Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_{W'})&=\mu_k\operatorname{sign}X\\&=\mu_k\operatorname{sign}W-\mu_k\operatorname{sign}W'.\end{aligned}$$

□

Here is the main theorem.

Theorem 3.32. *For $k\geq 1$, there is a constant $\mu_k\in\mathscr{A}_{2k,3k}$ such that*

$$\widehat{Z}_{2k,3k}(\vec{f})=Z_{2k,3k}(\vec{f})-Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W)+\mu_k\operatorname{sign}W\in\mathscr{A}_{2k,3k}$$

does not depend on the (generic) choices made. Thus $\widehat{Z}_{2k,3k}(\vec{f})$ is an invariant of diffeomorphism type of M.



FIGURE 5. Cerf's graphic

Proof of the theorem is given in §5. Theorem 3.32 allows us to put $\widehat{Z}_{2k,3k}(M) = \widehat{Z}_{2k,3k}(\vec{f})$.

Conjecture 3.33. *$\widehat{Z}_{2k,3k}(M)$ is nontrivial. In particular, the sequence $\{\widehat{Z}_{2k,3k}(M)\}_k$ recovers all Ohtsuki's finite type invariants (*[Oh96]*) over $\mathbb{Q}$.*

Conjecture 3.34. *$\widehat{Z}_{2k,3k}(M)$ agrees with the configuration space integral invariant of Axelrod–Singer (*[AS]*) and Kontsevich (*[Ko]*).*

4. **Moduli space of gradient flow graphs in 1-parameter family**

4.1. **Bifurcations in 1-parameter family.** We shall extend the moduli spaces $\mathscr{M}_2(f)$ and $\mathscr{M}_\Gamma(\vec{f})$ to those for 1-parameter families of functions $f_s : M \to \mathbb{R}$, $s \in [0,1]$ and metrics μ_s. To connect two Morse functions on a manifold in the space of smooth functions, 1-parameter families of Morse functions are not enough. The following fact is known.

Lemma 4.1 ([Ce])**.** *Two Morse functions on a manifold can be connected by a 1-parameter family of smooth functions with only Morse and birth/death singularities.*

Thus, the following bifurcations, that may change the topology of moduli spaces, may happen finitely many times in a generic 1-parameter family.

1. Level exchange point, i.e., the point where the order of the critical values changes.
2. Birth/death point.
3. Gradient i/i-intersection point, i.e., a point where a descending manifold and an ascending manifold of the same index i intersect transversally.

We may assume that no two different bifurcations overlap on a single time. In the following, we will call such a 1-parameter family a *generic 1-parameter family.*

It is convenient to represent bifurcations in a 1-parameter family by the graph of critical values, equipped with the information of gradient i/i-intersections. See Figure 5 for an example. Such a diagram is called a *graphic* ([Ce]). In a graphic, a level exchange point corresponds to a crossing of two curves, a gradient i/i-intersection between a pair of critical points is represented by a dotted arrow, and a birth/death point corresponds to beaks.

4.2. **Acyclicity in 1-parameter family.** When we associated an acyclic chain complex $\widetilde{C}^{(i)}$ to a Morse function $f_i : M \to \mathbb{R}$ in §3.1, we have chosen a pair $(a_i, b_i) \in P_d^{(i)} \times P_0^{(i)}$ of critical points. We show that two such choices can be connected by a pair of connected critical point loci in a generic 1-parameter family of functions.

Let $f_0, f_1 : M \to \mathbb{R}$ be two C^r Morse functions and take pairs $(a_0, b_0) \in P_d^{(0)} \times P_0^{(0)}$, $(a_1, b_1) \in P_d^{(1)} \times P_0^{(1)}$ of critical points of f_0, f_1 respectively.

Lemma 4.2. *There is a generic 1-parameter family of C^r functions $\{f_s\}_{s\in[0,1]}$ on M, considered as a single C^r-map $M \times [0,1] \to \mathbb{R}$, such that*

1. *$f_s : M \times \{s\} \to \mathbb{R}$ has only Morse or birth/death singularities for each s,*
2. *there are critical point loci $\widetilde{a}(s)$ and $\widetilde{b}(s)$ in $M \times [0,1]$ for $\{f_s\}$ that are nondegenerate on every fiber $M \times \{s\}$, with $\mathrm{pr}_1 \circ \widetilde{a}(0) = a_0$, $\mathrm{pr}_1 \circ \widetilde{a}(1) = a_1$, $\mathrm{pr}_1 \circ \widetilde{b}(0) = b_0$, $\mathrm{pr}_1 \circ \widetilde{b}(1) = b_1$.*

Thus we may assume the acyclicity through a generic 1-parameter family.

Proof. One can raise the level of a neighborhood of a_0 freely so that the level of a_0 is higher than other critical values of f_0. Similarly, one can lower the level of a neighborhood of b_0 so that the level of b_0 is lower than other critical values of f_0. For a small number $\varepsilon > 0$, we define f_s in $0 \leq s \leq \varepsilon$ as a 1-parameter family that gives the rising of a_0 and the falling of b_0.

We also do the same for (a_1, b_1), i.e., we define f_s in $1-\varepsilon \leq s \leq 1$ as a 1-parameter family that gives the rising of a_1 and the falling of b_1 as s decreases. Here we may assume that $f_\varepsilon(a_0) = f_{1-\varepsilon}(a_1)$ and $f_\varepsilon(b_0) = f_{1-\varepsilon}(b_1)$. We put $c_a = f_\varepsilon(a_0)$, $c_b = f_\varepsilon(b_0)$, $M' = f_\varepsilon^{-1}[c_b + \varepsilon', c_a - \varepsilon']$ and $M'' = f_{1-\varepsilon}[c_b + \varepsilon', c_a - \varepsilon']$ for a small number $\varepsilon' > 0$. Take smooth sections $\gamma_a, \gamma_b : [\varepsilon, 1-\varepsilon] \to M \times [\varepsilon, 1-\varepsilon]$ with disjoint images and their fiberwise tubular neighborhoods $N(\gamma_a)$ and $N(\gamma_b)$, so that

$$\begin{aligned}(N(\gamma_a) \textstyle\coprod N(\gamma_b)) \cap (M \times \{\varepsilon\}) &= M - M',\\ (N(\gamma_a) \textstyle\coprod N(\gamma_b)) \cap (M \times \{1-\varepsilon\}) &= M - M''.\end{aligned}$$

Then one can find a family of C^r functions

$$f'_s : (M \times \{s\}) \setminus (N(\gamma_a) \textstyle\coprod N(\gamma_b)) \to \mathbb{R},\ s \in [\varepsilon, 1-\varepsilon]$$

such that

1. $f'_\varepsilon = f_\varepsilon|_{M'}$ and $f'_{1-\varepsilon} = f_{1-\varepsilon}|_{M''}$,
2. f'_s is nonsingular on the boundary,
3. f'_s takes the maximum value $c_a - \varepsilon'$ on $\partial_+(M \times \{s\}) := \partial\overline{N(\gamma_a)} \cap (M \times \{s\})$ and $f'^{-1}_s(c_a - \varepsilon') = \partial_+(M \times \{s\})$,
4. f'_s takes the minimum value $c_b + \varepsilon'$ on $\partial_-(M \times \{s\}) := \partial\overline{N(\gamma_b)} \cap (M \times \{s\})$ and $f'^{-1}_s(c_b + \varepsilon') = \partial_-(M \times \{s\})$,
5. f'_s has only Morse or birth/death singularities.

Now the existence of the extension of f'_s on $N(\gamma_a) \coprod N(\gamma_n)$ so that the result is as desired, is straightforward. □

4.3. **Moduli space $\mathscr{M}_\Gamma$ in 1-parameter family and transversality.** Let $J = [s_0, s_1]$ be a closed interval in $[0,1]$. For a generic 1-parameter family $\vec{f_s} = (f_s, f_2, \ldots, f_m)$, $s \in J$, the moduli space $\mathscr{M}_\Gamma(\vec{f_s})$ is defined similarly as $\mathscr{M}_\Gamma(\vec{f})$ by replacing f_1 in the definition of $\mathscr{M}_\Gamma(\vec{f})$ with f_s if s is generic. Note that if $\beta(1) \in \mathrm{Se}(\Gamma)$, then $\mathscr{M}_\Gamma(\vec{f_s})$ is defined only on the complement of birth/death parameters in J, since the set of critical points changes at birth/death parameters. For graphs Γ with the dimension of the moduli space being negative (see the formula of Proposition 3.4), the moduli space $\mathscr{M}_\Gamma(\vec{f}_s)$ is empty at a generic parameter s, but we will see that $\mathscr{M}_\Gamma(\vec{f}_s)$ may be non-empty at finitely many non-generic parameters in J if $\dim \mathscr{M}_\Gamma(\vec{f}_s) = -1$.

We check the transversality on the moduli space $\mathscr{M}_\Gamma(\vec{f}_s)$, considered as the intersection, only for the graph Γ of Example 3.2 since other cases are similar. We fix the underlying sequence $\vec{C}$ of acyclic complexes for the graph as the restriction at $s = s_0$ of the family of acyclic Morse complexes obtained in §4.2. Suppose that $J = [s_0, s_1]$ does not include birth/death parameters and that $\vec{f}_{s_0} = (f_{s_0}, f_2, \ldots, f_6) \in (C^r(M))^6$ is generic in the sense of Proposition 3.4. We decompose Γ into two parts:

$\Gamma' =$ 

, and $\Gamma'' =$ 

.

The moduli space $\mathscr{M}_{\Gamma''}(\vec{f''})$, $\vec{f''} = (f_2, \ldots, f_6)$, is given by

$$\mathscr{M}_{\Gamma''}(\vec{f''}) = \mathrm{pr}_1 \circ \Phi_{\vec{f''}}^{-1}(\Delta''),$$

where $\Phi_{\vec{f''}} : \mathrm{Conf}_4(M) \times \mathbb{R}_+^5 \to M^9$ is defined by

$$\begin{aligned}&\Phi_{\vec{f''}}(x_1, x_2, x_3, x_4, t_2, \ldots, t_6)\\ &\quad = x_1 \times (x_2, \Phi_{f_2}^{t_2}(x_1)) \times (x_3, \Phi_{f_4}^{t_4}(x_1), \Phi_{f_5}^{t_5}(x_2)) \times (x_4, \Phi_{f_3}^{t_3}(x_2), \Phi_{f_6}^{t_6}(x_3)),\end{aligned}$$

and $\Delta'' = \{x_1 \times (x_2, x_2) \times (x_3, x_3, x_3) \times (x_4, x_4, x_4)\,;\, x_1, x_2, x_3, x_4 \in M\}$. The genericity of $\vec{f}_{s_0}$ implies that $\mathscr{M}_{\Gamma''}(\vec{f''})$ is a smooth submanifold of $\mathrm{Conf}_4(M)$ of dimension $(4d + 4) + 4d - 9d = 5 - d$. On the other hand, the moduli space $\mathscr{M}_{\Gamma'}(f_J) = \bigcup_{s \in J} \mathscr{M}_{\Gamma'}(f_s)$ is given by the $(d + \eta_1 + 1)$-dimensional manifold

$$\mathscr{M}_{\Gamma'}(f_J) = \bigcup_{s \in J} (\mathscr{D}_p(f_s) \times \mathscr{A}_q(f_s)) \subset \mathrm{Conf}_2(M) \times J.$$

Then we have

$$\mathscr{M}_\Gamma(\vec{f}_J) = \widetilde{\pi}_{14}^{-1}(\mathscr{M}_{\Gamma'}(f_s)) \cap (\mathscr{M}_{\Gamma''}(\vec{f''}) \times J),$$

where $\widetilde{\pi}_{14} : \mathrm{Conf}_4(M) \times J \to \mathrm{Conf}_2(M) \times J$ is the projection defined by $(x_1, x_2, x_3, x_4, s) \mapsto (x_1, x_4, s)$. By the transversality theorem, we may assume after a suitable small deformation of the family $\{(f_s, \mu_s)\}_{s \in J}$ that the intersection is transversal, and hence $\mathscr{M}_\Gamma(\vec{f}_J) = \bigcup_{s \in J} \mathscr{M}_\Gamma(\vec{f}_s)$ is a smooth submanifold of dimension

$$(d + \eta_1 + 1 + 2d) + (5 - d + 1) - (4d + 1) = 6 + \eta_1 - 2d.$$

The deformation can be chosen so small that it does not affect the genericity of $\vec{f}_0$. More generally, we have the following proposition.

Proposition 4.3. *Suppose that* $\Gamma \in \mathscr{G}_{n,m,\vec{\eta}}(\vec{C})$ *has no bivalent vertex. For a generic choice of a 1-parameter family* $\{(\vec{f}_s, \vec{\mu}_s)\}_{s \in J}$*, the space* $\mathscr{M}_\Gamma(\vec{f}_J)$ *is an oriented smooth manifold of dimension*

$$\dim \mathscr{M}_\Gamma(\vec{f}_J) = d(n - m) + \sum_{i=1}^{m} \eta_i + 1.$$

4.4. **Compactification of the moduli space of trajectories in 1-parameter family of Morse functions.** We construct a compactification of the moduli space $\mathscr{M}_2(f_J) = \bigcup_{s\in J} \mathscr{M}_2(f_s)$, where $J \subset [0,1]$ is an open interval that does not include birth/death parameters, i.e., f_J is a 1-parameter family of Morse functions.

4.4.1. *Moduli space on a level exchange point.* In the construction of $\overline{\mathscr{M}}_2(f)$ (in §3.3.2), we assumed that the critical values of f are all distinct. However, in 1-parameter family, the loci of critical values can not in general be ordered because of crossings of curves in a graphic. Instead, we consider the moduli space of "semi-short" trajectories that are close to an exchanging pair of critical point loci.

Let $u \in J$ be a point on which a level exchange bifurcation occurs and choose a small neighborhood J_u of u in J so that there are no other bifurcations on J_u. We shall construct a compactification of $\mathscr{M}_2(f_{J_u})$. Let $p = p_s, q = q_s$ be the pair of critical point loci of $f_{J_u} = \{f_s\}_{s\in J_u}$ that are in a level exchange position. Then there exists a smooth functions $\gamma_a, \gamma_b : J_u \to \mathbb{R}$ such that

1. $f_s(p_s), f_s(q_s) \in (\gamma_a(s), \gamma_b(s))$ for all $s \in J_u$,
2. for each $s \in J_u$, there are no critical points of f_s in $f_s^{-1}[\gamma_a(s), \gamma_b(s)]$ except p_s and q_s.

We put $\widetilde{L}_a = \bigcup_{s\in J_u} f_s^{-1}(\gamma_a(s))$, $\widetilde{L}_b = \bigcup_{s\in J_u} f_s^{-1}(\gamma_b(s))$, $W_{pq}(s) = f_s^{-1}[\gamma_a(s), \gamma_b(s)]$, $\widetilde{W}_{pq} = \bigcup_{s\in J_u} W_{pq}(s)$. Let

$$\begin{aligned}\widetilde{K}_p &= \bigcup_{s\in J_u} (\mathscr{D}_{p_s}(f_s) \cup \mathscr{A}_{p_s}(f_s)) \cap W_{pq}(s),\\ \widetilde{K}_q &= \bigcup_{s\in J_u} (\mathscr{D}_{q_s}(f_s) \cup \mathscr{A}_{q_s}(f_s)) \cap W_{pq}(s).\end{aligned}$$

We take small neighborhoods $\widetilde{B}_p$ and $\widetilde{B}_q$ of $\widetilde{K}_p \cap \widetilde{L}_b$ and $\widetilde{K}_q \cap \widetilde{L}_b$ in $\widetilde{L}_b$ respectively. Let $\widetilde{A}_p \subset \widetilde{L}_a$ be the union of $\widetilde{K}_p \cap \widetilde{L}_a$ and the subset of $\widetilde{L}_a$ consisting of points (s, ℓ) such that $\ell = \Phi^t_{f_s}(x)$ for some $t > 0$ and for a point $x \in \widetilde{B}_p \cap L_p(s)$. In other words, $\widetilde{A}_p$ be the union of $\widetilde{K}_p \cap \widetilde{L}_a$ and the image of the negative gradient flow from $\widetilde{B}_p \cap \widetilde{L}_b$. The subset $\widetilde{A}_q$ of $\widetilde{L}_a$ is similarly defined.

Since $\widetilde{K}_p \cap \widetilde{K}_q = \emptyset$, we may assume that any trajectory starting from $\widetilde{B}_p$ are disjoint from trajectories starting from $\widetilde{B}_q$. Thus we have

$$\mathscr{M}_2(f_{J_u}; \widetilde{B}_p, \widetilde{L}_a) = \mathscr{M}_2(f_{J_u}; \widetilde{B}_p, \widetilde{A}_p), \quad \mathscr{M}_2(f_{J_u}; \widetilde{B}_q, \widetilde{L}_a) = \mathscr{M}_2(f_{J_u}; \widetilde{B}_q, \widetilde{A}_q)$$

and in particular, the two moduli spaces are disjoint as subspaces of $\mathscr{M}_2(f_{J_u}; \widetilde{L}_b, \widetilde{L}_a)$. Since each of $\mathscr{M}_2(f_{J_u}; \widetilde{B}_p, \widetilde{A}_p)$ and $\mathscr{M}_2(f_{J_u}; \widetilde{B}_p, \widetilde{A}_p)$ has only one end for broken trajectories, the compactifications

$$\overline{\mathscr{M}}_2(f_{J_u}; \widetilde{B}_p, \widetilde{A}_p), \quad \overline{\mathscr{M}}_2(f_{J_u}; \widetilde{B}_q, \widetilde{A}_q)$$

can be defined in a similar way as §3.3 (using parameterized Morse lemma [Ig2, Appendix]). So it may be natural to define

$$\begin{aligned}\overline{\mathscr{M}}_2(f_{J_u}; \widetilde{B}_p, \widetilde{L}_a) &= \overline{\mathscr{M}}_2(f_{J_u}; \widetilde{B}_p, \widetilde{A}_p),\\ \overline{\mathscr{M}}_2(f_{J_u}; \widetilde{B}_q, \widetilde{L}_a) &= \overline{\mathscr{M}}_2(f_{J_u}; \widetilde{B}_q, \widetilde{A}_q).\end{aligned}$$

Then we construct an extension of $\overline{\mathscr{M}}_2(f_{J_u};\widetilde{B}_p,\widetilde{L}_a)\coprod\overline{\mathscr{M}}_2(f_{J_u};\widetilde{B}_q,\widetilde{L}_a)$ to $\overline{\mathscr{M}}_2(f_{J_u};\widetilde{L}_b,\widetilde{L}_a)$ as follows. Let $\widetilde{X}\subset\widetilde{L}_b$ be an open subset such that

(1) $\widetilde{X}\cup\widetilde{B}_p\cup\widetilde{B}_q=\widetilde{L}_b$,
(2) $\widetilde{X}\cap\widetilde{K}_p=\emptyset$, $\widetilde{X}\cap\widetilde{K}_q=\emptyset$.

Then the negative gradient flow carries $\widetilde{X}$ diffeomorphically onto an open subset $\widetilde{Y}$ of $\widetilde{L}_a$. Obviously, $\widetilde{Y}$ satisfies the following conditions:

(1) $\widetilde{Y}\cup\widetilde{A}_p\cup\widetilde{A}_q=\widetilde{L}_a$,
(2) $\widetilde{Y}\cap\widetilde{K}_p=\emptyset$, $\widetilde{X}\cap\widetilde{K}_q=\emptyset$.

Since there are no ends in $\mathscr{M}_2(f_{J_u};\widetilde{X},\widetilde{Y})$ for broken trajectories, we define

$$\overline{\mathscr{M}}_2(f_{J_u};\widetilde{X},\widetilde{L}_b)=\mathscr{M}_2(f_{J_u};\widetilde{X},\widetilde{L}_b)=\mathscr{M}_2(f_{J_u};\widetilde{X},\widetilde{Y}).$$

Now we define

$$\overline{\mathscr{M}}_2(f_{J_u};\widetilde{L}_b,\widetilde{L}_a)=\overline{\mathscr{M}}_2(f_{J_u};\widetilde{B}_p,\widetilde{L}_a)\cup\overline{\mathscr{M}}_2(f_{J_u};\widetilde{X},\widetilde{L}_a)\cup\overline{\mathscr{M}}_2(f_{J_u};\widetilde{B}_q,\widetilde{L}_a),$$

where they are glued along $\mathscr{M}_2(f_{J_u};\widetilde{B}_p\cap\widetilde{X},\widetilde{L}_a)$ and $\mathscr{M}_2(f_{J_u};\widetilde{B}_q\cap\widetilde{X},\widetilde{L}_a)$ via the inclusions. The compactifications $\overline{\mathscr{M}}_2(f_{J_u};\widetilde{L}_b,\widetilde{W}_{pq})$, $\overline{\mathscr{M}}_2(f_{J_u};\widetilde{W}_{pq},\widetilde{L}_a)$ etc. can be constructed by a similar gluing construction from moduli spaces of trajectories around a single critical point locus.

Over the interval J_u, we may assume that critical point loci except p and q are ordered, and after a suitable small deformation of the family of Morse pairs that does not change the graphic, we may assume that (f_s,μ_s) is Morse–Smale for all $s\in J_s$. So similar extensions as in §3.3 works for other critical point loci and we will finally get a compactification

$$\overline{\mathscr{M}}_2(f_{J_u})=\overline{\mathscr{M}}_2(f_{J_u};J_u\times M,J_u\times M).$$

By the same construction at all the level exchange points $u_1,u_2,\ldots,u_r\in J_u$, we will obtain a compactification on $\coprod_{j=1}^r J_{u_j}$.

4.4.2. *Moduli space on ordered 1-parameter family.* We would like to extend the compactifications of moduli spaces on $\coprod_{j=1}^r J_{u_j}$, obtained in §4.4.1, over the whole of J. We assume $u_1<u_2<\cdots<u_r$. Let $I_j\subset J$, $j=0,1,2,\ldots,r$ be a sequence of mutually disjoint open intervals such that

(1) $\bigcup_{j=0}^r I_j\cup\bigcup_{j=1}^r J_{u_j}=J$,
(2) $(\coprod_{j=0}^r I_j)\cap\{u_1,\ldots,u_r\}=\emptyset$,
(3) $I_j\subset(u_j,u_{j+1})$.

See Figure 6. We shall construct a compactification $\overline{\mathscr{M}}_2(f_{I_j})$ of $\mathscr{M}_2(f_{I_j})$ connecting $\overline{\mathscr{M}}_2(f_{J_{u_j}})$ and $\overline{\mathscr{M}}_2(f_{J_{u_{j+1}}})$.

Over each I_j, the critical values are ordered everywhere, so we can separate critical point loci into small pieces by families of level surfaces. We already have a compactification on each piece which includes only one critical point locus. However, there is a problem in making the intersection between two moduli spaces as in Lemma 3.8, because an analogue of the Morse–Smale condition for 1-parameter family, which requires the Morse–Smale condition on each fiber, fails. Instead, we consider as follows: Although we can not require fiberwise Morse–Smale condition, we can require the transversality between a family of descending manifolds and a



FIGURE 6.

family of ascending manifolds. This requirement suffices for the moduli space to be a smooth submanifold of a fiber bundle over I_j (with fiber $\mathrm{Conf}_2(M)$), though it may not be a subbundle. See §3.1 of [Wa] for detail.

Let $\widetilde{\mathscr{A}_p}(f_{I_j}) = \bigcup_{s\in I_j}\mathscr{A}_{p_s}(f_s)$ and $\widetilde{\mathscr{D}_p}(f_{I_j}) = \bigcup_{s\in I_j}\mathscr{D}_{p_s}(f_s)$. By changing the 1-parameter family $\{\mu_s\}_{s\in I_j}$ of metrics on M over I_j, one can show by a similar argument as the proof of the genericity of Morse–Smale condition, that generically $\widetilde{\mathscr{A}_p}(f_{I_j})$'s and $\widetilde{\mathscr{D}_q}(f_{I_j})$'s intersect mutually transversal in the trivial M-bundle over I_j. Thus we will get the following proposition, which is an analogue of Proposition 3.10.

Proposition 4.4. *For a generic 1-parameter family, $\mathscr{M}_2(f_{I_j})$ has a canonical compactification $\overline{\mathscr{M}}_2(f_{I_j})$ to a smooth manifold with corners, whose codimension k stratum consists of families of k times broken trajectories. Hence we have a canonical compactification $\overline{\mathscr{M}}_2(f_J)$ of $\mathscr{M}_2(f_J)$ over J.*

By using the compactification $\overline{\mathscr{M}}_2(f_J)$, one can also define the compactification $\overline{\mathscr{M}}_\Gamma(f_J)$ of $\mathscr{M}_\Gamma(f_J)$. Then we have the following corollary.

Corollary 4.5. *If Γ does not have bivalent vertices, then on a neighborhood of the strata of codimension ≤ 1, $\overline{\mathscr{M}}_\Gamma(f_J)$ is a smooth manifold with boundary, that is a smooth submanifold of $J \times \overline{\mathrm{Conf}}_n(M)$ with boundary. The boundary consists of flows with a broken trajectory or with a subgraph collapsed.*

4.5. **Gluing at birth/death bifurcation.** Let $s_0 \in [0,1]$ be a death parameter in a generic 1-parameter family and let $J_{s_0} \subset [0,1]$ be a small open interval around s_0. Let $v \in M$ be the death point at s_0. By the parametrized normal form lemma for birth/death singularity (e.g., [Ig2, Appendix], [Ce]), there is a local coordinate on a neighborhood M_v of v in $J_{s_0} \times M$ on which f_s agrees with

$$h_u(x) = c(u) + \frac{x_1^3}{3} + ux_1 - \frac{x_2^2}{2} - \cdots - \frac{x_i^2}{2} + \frac{x_{i+1}^2}{2} + \cdots + \frac{x_d^2}{2}, \quad u \in \mathbb{R},$$

where u is a rescaling of s and $c(u)$ is a smooth function of u, and one can deform the metric on M_v so that it agrees on M_v with the restriction of the standard metric on $\mathbb{R}^d$. The negative gradient field of h_u with respect to the standard metric is

$$-\mathrm{grad}\, h_u = (-x_1^2 - u, x_2, \ldots, x_i, -x_{i+1}, \ldots, -x_d).$$

So, on $u > 0$, there are no critical points of h_u. At $u = 0$, there is only one critical point at the origin, and on $u < 0$, there are exactly two critical points

$p_\pm = (\pm\sqrt{u}, 0, \ldots, 0)$ of h_u. We shall describe how a pair of trajectories going from/to critical points of h_u on $u < 0$ are glued together into a single trajectory on $u > 0$.

4.5.1. *Gradient trajectories of h_u, $u > 0$ case.* The integral curve $\gamma : \mathbb{R} \to \mathbb{R}^d$, $\gamma(t) = (\gamma_1(t), \ldots, \gamma_d(t))$ of $-\mathrm{grad}\, h_u$ is determined by the system of differential equations:

$$\begin{aligned} &\frac{d\gamma_1(t)}{dt} = -\gamma_1(t)^2 - u,\ \frac{d\gamma_2(t)}{dt} = \gamma_2(t),\ \ldots,\ \frac{d\gamma_i(t)}{dt} = \gamma_i(t),\\ &\frac{d\gamma_{i+1}(t)}{dt} = -\gamma_{i+1}(t),\ \ldots,\ \frac{d\gamma_d(t)}{dt} = -\gamma_d(t), \end{aligned} \tag{4.1}$$

for each given initial point $(\gamma_1(0), \ldots, \gamma_d(0))$. In $u > 0$, the solution of (4.1) is given by

$$\begin{aligned} &\gamma_1(t) = \frac{\sqrt{u}\,\gamma_1(0) - u\tan\sqrt{u}\,t}{\sqrt{u} + \gamma_1(0)\tan\sqrt{u}\,t},\ \gamma_2(t) = \gamma_2(0)e^t,\ \ldots,\ \gamma_i(t) = \gamma_i(0)e^t,\\ &\gamma_{i+1}(t) = \gamma_{i+1}(0)e^{-t},\ \ldots,\ \gamma_d(t) = \gamma_d(0)e^{-t}. \end{aligned}$$

For a small number $\varepsilon > 0$, let L_ε and $L_{-\varepsilon}$ be the subsets of $\mathbb{R}^d$ given by

$$\begin{aligned} L_\varepsilon &= \{(\varepsilon, x_2, \ldots, x_d)\,;\, x_2, \ldots, x_d \in \mathbb{R}\},\\ L_{-\varepsilon} &= \{(-\varepsilon, x_2, \ldots, x_d)\,;\, x_2, \ldots, x_d \in \mathbb{R}\}. \end{aligned}$$

Since $-\mathrm{grad}_{(0,x_2,\ldots,x_d)}\, h_u = (-u, x_2, \ldots, x_i, -x_{i+1}, \ldots, -x_d)$, one may see that any trajectory of h_u in $u > 0$ and in M_v intersects both L_ε and $L_{-\varepsilon}$. Conversely, for any point a of $L_\varepsilon \cap M_v$ (resp. $L_{-\varepsilon} \cap M_v$), there exists a unique (shift equivalence class of) gradient trajectory of h_u which intersects $L_\varepsilon \cap M_v$ (resp. $L_{-\varepsilon} \cap M_v$) at a. So there is a one-to-one correspondence between a point on L_ε or $L_{-\varepsilon}$ and a gradient trajectory of h_u.

Now suppose that an integral curve $\gamma(t)$ of $-\mathrm{grad}\, h_u$ starts at a point of L_ε. We shall describe the point $\mathrm{Im}\,\gamma \cap L_{-\varepsilon}$. If $\gamma(t_{-\varepsilon}) \in L_{-\varepsilon}$ at $t_{-\varepsilon} > 0$, then by (4.1),

$$t_{-\varepsilon} = -\int_\varepsilon^{-\varepsilon} \frac{dx}{x^2 + u} = -\frac{2}{\sqrt{u}}\mathrm{Tan}^{-1}\frac{\varepsilon}{\sqrt{u}} = -\frac{1}{\sqrt{u}}\Bigl(\mathrm{Tan}^{-1}\frac{2\varepsilon\sqrt{u}}{u - \varepsilon^2} + \pi\Bigr)$$

for $0 < u < \varepsilon^2$. We put $\tau_\varepsilon(u) = \frac{1}{\sqrt{u}}\Bigl(\mathrm{Tan}^{-1}\frac{2\varepsilon\sqrt{u}}{u-\varepsilon^2} + \pi\Bigr)$ for simplicity. Then $\tau_\varepsilon(u)$ has the following expansion (convergent on $0 < u < \varepsilon^2$):

$$\tau_\varepsilon(u) = \frac{\pi}{\sqrt{u}} - 2\sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)\varepsilon^{2k+1}} u^k. \tag{4.2}$$

The point $\gamma(t_{-\varepsilon})$ can be expressed by using $\tau_\varepsilon(u)$ as follows.

$$\gamma(t_{-\varepsilon}) = \bigl(-\varepsilon, \gamma_2(0)e^{\tau_\varepsilon(u)}, \ldots, \gamma_i(0)e^{\tau_\varepsilon(u)}, \gamma_{i+1}(0)e^{-\tau_\varepsilon(u)}, \ldots, \gamma_d(0)e^{-\tau_\varepsilon(u)}\bigr).$$

If we put $\varepsilon_2 = \gamma_2(0)e^{\tau_\varepsilon(u)}, \ldots, \varepsilon_i = \gamma_i(0)e^{\tau_\varepsilon(u)}, \varepsilon_{i+1} = \gamma_{i+1}(0), \ldots, \varepsilon_d = \gamma_d(0)$, then the integral curve starting at the point

$$\bigl(\varepsilon, \varepsilon_2 e^{-\tau_\varepsilon(u)}, \ldots, \varepsilon_i e^{-\tau_\varepsilon(u)}, \varepsilon_{i+1}, \ldots, \varepsilon_d\bigr) \in L_\varepsilon \tag{4.3}$$

intersects $L_{-\varepsilon}$ at the point

$$\bigl(-\varepsilon, \varepsilon_2, \ldots, \varepsilon_i, \varepsilon_{i+1}e^{-\tau_\varepsilon(u)}, \ldots, \varepsilon_d e^{-\tau_\varepsilon(u)}\bigr) \in L_{-\varepsilon}. \tag{4.4}$$



FIGURE 7.

This observation suggests a way to glue trajectories.

4.5.2. *Gradient trajectories of h_u going from/to critical points, $u \leq 0$ case.* In $u < 0$, the ascending/descending manifolds of h_u are described as follows. See Figure 7.

$$\mathscr{A}_{p_+}(h_u) = \{x_2 = \cdots = x_i = 0, x_1 \geq -\sqrt{u}\},\ \mathscr{D}_{p_+}(h_u) = \{x_{i+1} = \cdots = x_d = 0, x_1 = \sqrt{u}\},$$
$$\mathscr{A}_{p_-}(h_u) = \{x_2 = \cdots = x_i = 0, x_1 = -\sqrt{u}\},\ \mathscr{D}_{p_-}(h_u) = \{x_{i+1} = \cdots = x_d = 0, x_1 \leq \sqrt{u}\}.$$

Hence

$$\begin{aligned} L_\varepsilon \cap \mathscr{A}_{p_+}(h_u) &= \{(\varepsilon, 0, \ldots, 0, \varepsilon_{i+1}, \ldots, \varepsilon_d) \in \mathbb{R}^d\,;\, \varepsilon_{i+1}, \ldots, \varepsilon_d \in \mathbb{R}\}, \\ L_{-\varepsilon} \cap \mathscr{D}_{p_-}(h_u) &= \{(-\varepsilon, \varepsilon_2, \ldots, \varepsilon_i, 0, \ldots, 0) \in \mathbb{R}^d\,;\, \varepsilon_2, \ldots, \varepsilon_i \in \mathbb{R}\}. \end{aligned} \tag{4.5}$$

One may check that this holds also for $u = 0$, in which case $p_+ = p_- = v$.

4.5.3. *Gluing at $u = 0$.* We define a map

$$\varphi : (-\varepsilon^2, \varepsilon^2) \times \mathbb{R}^{d-1} \to (-\varepsilon^2, \varepsilon^2) \times \mathbb{R}^{d-1} \times \mathbb{R}^{d-1}$$

by

$$\varphi(u; \varepsilon_2, \ldots, \varepsilon_d) = \begin{cases} u \times \big(\varepsilon_2 e^{-\tau_\varepsilon(u)}, \ldots, \varepsilon_i e^{-\tau_\varepsilon(u)}, \varepsilon_{i+1}, \ldots, \varepsilon_d\big) & \\ \quad \times \big(\varepsilon_2, \ldots, \varepsilon_i, \varepsilon_{i+1} e^{-\tau_\varepsilon(u)}, \ldots, \varepsilon_d e^{-\tau_\varepsilon(u)}\big), & \text{if } u > 0 \\ u \times \big(0, \ldots, 0, \varepsilon_{i+1}, \ldots, \varepsilon_d\big) \times \big(\varepsilon_2, \ldots, \varepsilon_i, 0, \ldots, 0\big), & \text{if } u \leq 0 \end{cases}$$

We consider the space of gradient trajectories of h_u or pairs of gradient trajectories of h_u intersecting both L_ε and $L_{-\varepsilon}$, as a subspace of $L_\varepsilon \times L_{-\varepsilon}$. By (4.5), the part $\varphi((-\varepsilon^2, 0] \times \mathbb{R}^{d-1})$ is the space of pairs $(\gamma_{p_+}(t), \gamma_{p_-}(t))$ of integral curves of $-\mathrm{grad}\, h_u$ satisfying the conditions

$$\begin{aligned} \lim_{t \to \infty} \gamma_{p_+}(t) = p_+,\ \gamma_{p_+}(0) \in L_\varepsilon, \\ \lim_{t \to -\infty} \gamma_{p_-}(t) = p_-,\ \gamma_{p_-}(0) \in L_{-\varepsilon}. \end{aligned} \tag{4.6}$$

On the other hand, by (4.3) and (4.4), the part $\varphi((0, \varepsilon^2) \times \mathbb{R}^{d-1})$ is the space of negative gradient trajectories of h_u, $u > 0$, near the origin.



FIGURE 8. The image of φ (in the $u\varepsilon_2\varepsilon_2'$-space). Example of $d = 2$ case : $\varphi(u; \varepsilon_2) = u \times (\varepsilon, \varepsilon_2) \times (-\varepsilon, \varepsilon_2 e^{-\tau_\varepsilon(u)})$.



FIGURE 9. Gluing of trajectories. $d = 3$ case.

Proposition 4.6. *The map φ is smooth and is an embedding. Hence any smooth path ξ on $\operatorname{Im}\varphi$ that goes from $u < 0$ to $u = 0$ with non-degenerate tangent on $T_0(-\varepsilon^2, \varepsilon^2)$ has a smooth extension on $u > 0$ part. In other words, a pair $(\gamma_{p_+}(t), \gamma_{p_-}(t))$ of integral curves of $-\operatorname{grad} h_u$ as (4.6) above are glued together at $u = 0$ into a single trajectory if the pair goes along the path ξ (as in Figure 9).*

For the proof of Proposition 4.6, we need two lemmas.

Lemma 4.7. *There exist series $P_n(u), Q_n(u) \in \mathbb{R}[[u]][\sqrt{u}]$, $n \geq 0$, that are well-defined as C^∞ differentiable functions on $(0, \varepsilon^2)$ such that*

$$\frac{d^n}{du^n} e^{-\tau_\varepsilon(u)} = \frac{Q_n(u) + \pi^n \varepsilon^{2n}}{P_n(u)} e^{-\tau_\varepsilon(u)},$$
$$\lim_{u \to 0} P_n(u) = \lim_{u \to 0} Q_n(u) = 0.$$

Proof. We prove the lemma by induction on n. The case $n = 0$ is obvious. Suppose the assertion holds true for n. Then we have

$$\begin{aligned}
\frac{d^{n+1}}{du^{n+1}} e^{-\tau_\varepsilon(u)} &= \frac{d}{du}\left(\frac{Q_n(u)+\pi^n\varepsilon^{2n}}{P_n(u)} e^{-\tau_\varepsilon(u)}\right) \\
= -\,&\frac{e^{-\tau_\varepsilon(u)}}{\sqrt{u}(2u^2+2\varepsilon^2 u)P_n(u)} \\
&\times\Bigg[\sqrt{u}\Big\{\Big((2u+2\varepsilon^2)\frac{uP_n'(u)}{P_n(u)}-2\varepsilon\Big)\pi^n\varepsilon^{2n}+(-2u-2\varepsilon^2)uQ_n'(u) \\
&\qquad + (2u+2\varepsilon^2)Q_n(u)\frac{uP_n'(u)}{P_n(u)} - 2\varepsilon Q_n(u)\Big\} \\
&\quad + \Big\{(-u-\varepsilon^2)\mathrm{Tan}^{-1}\frac{2\varepsilon\sqrt{u}}{u-\varepsilon^2}+(-\pi u-\pi\varepsilon^2)\Big\}\pi^n\varepsilon^{2n} \\
&\quad + (-u-\varepsilon^2)Q_n(u)\mathrm{Tan}^{-1}\frac{2\varepsilon\sqrt{u}}{u-\varepsilon^2}+(-\pi u-\pi\varepsilon^2)Q_n(u)\Bigg].
\end{aligned}\tag{4.7}$$

By (4.2), one may see that

$$\mathrm{Tan}^{-1}\frac{2\varepsilon\sqrt{u}}{u-\varepsilon^2} \in \mathbb{R}[[u]][\sqrt{u}], \quad \frac{uP_n'(u)}{P_n(u)} \in \mathbb{R}[[u]][\sqrt{u}],$$

$$\lim_{u\to 0}\mathrm{Tan}^{-1}\frac{2\varepsilon\sqrt{u}}{u-\varepsilon^2} = 0, \quad \lim_{u\to 0} uQ_n'(u) = 0, \quad \lim_{u\to 0}\frac{uP_n'(u)}{P_n(u)} < \infty.$$

This implies that the right hand side of (4.7) is of the form

$$\frac{Q_{n+1}(u)+\pi^{n+1}\varepsilon^{2n+2}}{\sqrt{u}(2u^2+2\varepsilon^2 u)P_n(u)} e^{-\tau_\varepsilon(u)}$$

for some C^∞ differentiable function $Q_{n+1}(u) \in \mathbb{R}[[u]][\sqrt{u}]$ with $\lim_{u\to 0} Q_{n+1}(u) = 0$ that is well-defined on $(0, \varepsilon^2)$. The proof completes if we put $P_{n+1}(u) = \sqrt{u}(2u^2 + 2\varepsilon^2 u)P_n(u)$. $\square$

Lemma 4.8. *For all $n \geq 0$,*

$$\lim_{u\to 0}\frac{d^n}{du^n}e^{-\tau_\varepsilon(u)} = 0.$$

Proof. Since $\lim_{u\to 0}(Q_n(u) + \pi^n\varepsilon^{2n}) = \pi^n\varepsilon^{2n}$, it suffices to show that

$$\lim_{u\to 0}\frac{e^{-\tau_\varepsilon(u)}}{P_n(u)} = \lim_{u\to 0}\exp(-\tau_\varepsilon(u) - \log P_n(u)) = 0.$$

The assertion follows if we show that $\lim_{u\to 0}(-\tau_\varepsilon(u) - \log P_n(u)) = -\infty$.

We put $s_n(u) = -\tau_\varepsilon(u) - \log P_n(u)$. Then we have

$$\begin{aligned}
\lim_{u\to 0} us_n'(u) &= \lim_{u\to 0}\Big(\frac{(u+\varepsilon^2)\mathrm{Tan}^{-1}\big(\frac{2\varepsilon\sqrt{u}}{u-\varepsilon^2}\big)+\pi u+2\varepsilon\sqrt{u}+\pi\varepsilon^2}{\sqrt{u}(2u+2\varepsilon^2)} - \frac{uP_n'(u)}{P_n(u)}\Big) \\
&= \infty
\end{aligned}$$

and hence $\lim_{u\to 0} s_n'(u) = \infty$. So we must have $\lim_{u\to 0} s_n(u) = -\infty$. $\square$

We put

$$\sigma_\varepsilon(u) = \begin{cases} e^{-\tau_\varepsilon(u)} & \text{if } u > 0 \\ 0 & \text{if } u \leq 0 \end{cases}$$

Then Lemma 4.8 shows that $\sigma_\varepsilon(u)$ is C^∞ differentiable.

Proof of Proposition 4.6. The map φ can be rewritten as

$$\begin{aligned}\varphi(u;\varepsilon_2,\ldots,\varepsilon_d) = u \times (&\varepsilon_2\sigma_\varepsilon(u),\ldots,\varepsilon_i\sigma_\varepsilon(u),\varepsilon_{i+1},\ldots,\varepsilon_d)\\ &\times(\varepsilon_2,\ldots,\varepsilon_i,\varepsilon_{i+1}\sigma_\varepsilon(u),\ldots,\varepsilon_d\sigma_\varepsilon(u)).\end{aligned}$$

This shows that φ is C^∞ differentiable. That the Jacobian matrix has full rank is obvious from the definition of φ on $u \leq 0$. Hence φ is an embedding. □

Recall that $s_0 \in [0,1]$ is a death parameter. Let (p_1, q_1) be the pair of critical points of $f_{s_0-\varepsilon'}$ for sufficiently small number $\varepsilon' > 0$, such that $d(p_1) = d(q_1) + 1 = i + 1$ and such that they are eliminated on $s > s_0$ after passing through the death point v. Let $\Gamma(p_1,q_1)_1 \in \mathscr{G}_{n,m}(\vec{C}_{s_0-\varepsilon'})$ be a graph with no bivalent vertices and let $\Gamma(\emptyset,\emptyset)_1 \in \mathscr{G}_{n,m}(\vec{C}_{s_0\pm\varepsilon'})$ be the graph obtained from $\Gamma(p_1,q_1)_1$ by replacing the edge $\beta(1)$ with a compact edge.

If $d = 3$ and if $3n = 2m$, then by Proposition 3.4, $\dim \overline{\mathscr{M}}_\Gamma(\vec{f_s}) = 0$ for $\Gamma \in \mathscr{G}_{n,m}(\vec{C})$. Since the moduli space is oriented, the number $\#\overline{\mathscr{M}}_\Gamma(\vec{f_s})$ can be counted with signs.

Corollary 4.9. *Suppose $d = 3$ and $3n = 2m$. The embedding φ induces a smooth cobordism between $\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_1}(\vec{f}_{s_0+\varepsilon'})$ and $-\overline{\mathscr{M}}_{\Gamma(p_1,q_1)_1}(\vec{f}_{s_0-\varepsilon'}) \coprod -\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_1}(\vec{f}_{s_0-\varepsilon'})$. Hence we have*

$$\#\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_1}(\vec{f}_{s_0+\varepsilon'}) = \#\overline{\mathscr{M}}_{\Gamma(p_1,q_1)_1}(\vec{f}_{s_0-\varepsilon'}) + \#\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_1}(\vec{f}_{s_0-\varepsilon'}).$$

4.6. **Orientation in 1-parameter family.** For a graph $\Gamma \in \mathscr{G}_{2k,3k}(\vec{C})$, we consider the orientation of $\partial\overline{\mathscr{M}}_\Gamma(\vec{f}_J)$ induced from $o^*(\overline{\mathscr{M}}_\Gamma(\vec{f}_J))$.

Choose a number k in $1 \leq k \leq m$ and put

$$\begin{aligned}&\widetilde{H}_j = J \times H_j,\ \widetilde{\Theta}_j = J \times \Theta_j \subset J \times \mathrm{Conf}_{2k}(M),\\ &\widetilde{\Sigma}_k = \begin{cases} \bigcap_{\substack{1\leq j\leq a\\ j\neq k}} \widetilde{H}_j \cap \bigcap_{j=a+1}^m \widetilde{\Theta}_j & \text{if } 1 \leq k \leq a\\ \bigcap_{j=1}^a \widetilde{H}_j \cap \bigcap_{\substack{a+1\leq j\leq m\\ j\neq k}} \widetilde{\Theta}_j & \text{if } a+1 \leq k \leq m\end{cases}\end{aligned}$$

Then

$$\begin{aligned}\mathrm{codim}\,\widetilde{\Sigma}_k &= \mathrm{codim}\,\mathscr{M}_\Gamma(\vec{f}_J) - \mathrm{codim}\,\widetilde{H}_k\\ &= (dn + 1 - d(n-m) - m - 1) - (d + \eta_k)\\ &= (d-1)(m+1) - 2d + 1 - \eta_k \equiv 0 \pmod 2\end{aligned}$$

Let X be a graph obtained from Γ with one edge replaced by a broken edge so that $\mathscr{M}_X(\vec{f}_J)$ is 0-dimensional. Suppose for simplicity that separated edges of Γ are labeled $1, 2, \ldots, a$. We shall describe the orientation of the face $\mathscr{S}_X$ of $\partial\overline{\mathscr{M}}_\Gamma(\vec{f}_J)$ corresponding to X, induced from the standard orientation of $\mathscr{M}_\Gamma(\vec{f}_J)$. We define the standard orientation of $\mathscr{M}_\Gamma(\vec{f}_J)$ by the formula

$$o^*(\mathscr{M}_\Gamma(\vec{f}_J)) = \bigwedge_{j=1}^a o^*(\widetilde{H}_j) \wedge \bigwedge_{j=a+1}^m o^*(\widetilde{\Theta}_j).$$

(1) $X =$ 

. For $\mathscr{M}_X(\vec{f}_J)$ to be 0-dimensional, $d(r_k) = d(p_k) - 1$ or $d(r_k) = d(p_k)$. The same statement as Lemma 3.15 holds true even if $\mathscr{D}_p(f)$ is replaced with a 1-parameter family of descending manifolds of p. The orientation of $\mathscr{F}_{r_k}\overline{\mathscr{M}}_\Gamma(\vec{f}_J)$ induced from the standard one

$$o^*(\mathscr{M}_\Gamma(\vec{f}_J)) = o^*(\widetilde{\mathscr{D}}_{p_k}(f_k)) \wedge o^*(\widetilde{\mathscr{A}_{q_k}}(f_k)) \wedge o^*(\widetilde{\Sigma}_k)$$

is given by

$$\begin{aligned}&o^*(\overline{\mathscr{M}(f_k;p_k,r_k)}) \wedge o^*(\widetilde{\mathscr{D}}_{r_k}(f_k)) \wedge o^*(\widetilde{\mathscr{A}_{q_k}}(f_k)) \wedge o^*(\widetilde{\Sigma}_k)\\ &= o^*(\overline{\mathscr{M}(f_k;p_k,r_k)}) \wedge o^*(\mathscr{M}_{\Gamma'}(\vec{f}_J)).\end{aligned}$$

This agrees with the standard orientation $o^*(\mathscr{M}_\Gamma(\vec{f}_J))$.

(2) $X =$ 

. For $\mathscr{M}_X(\vec{f}_J)$ to be 0-dimensional, $d(s_k) = d(q_k) + 1$ or $d(s_k) = d(q_k)$. If $\mathscr{A}_q(f)$ in the statement of Lemma 3.16 is replaced with a 1-parameter family of ascending manifolds of q, then the induced orientation would be

$$(-1)^{d(r)-(d(q)+1)} o^*(\widetilde{\mathscr{A}_r}(f))_x \wedge o^*(\overline{\mathscr{M}(f_k;r,q)})_{x'},$$

since the sign $(-1)^{k+1}$ in the formula (3.7) becomes $(-1)^{d(r)-(d(q)+1)}$. Thus, the orientation of $\mathscr{F}_{s_k}\overline{\mathscr{M}}_\Gamma(\vec{f}_J)$ induced from the standard one

$$o^*(\mathscr{M}_\Gamma(\vec{f}_J)) = o^*(\widetilde{\Sigma_k}) \wedge o^*(\widetilde{\mathscr{D}}_{p_k}(f_k)) \wedge o^*(\widetilde{\mathscr{A}_{q_k}}(f_k))$$

is given by

$$\begin{aligned}&(-1)^{d(s_k)-d(q_k)-1} o^*(\widetilde{\Sigma_k}) \wedge o^*(\widetilde{\mathscr{D}}_{p_k}(f_k)) \wedge o^*(\widetilde{\mathscr{A}_{s_k}}(f_k)) \wedge o^*(\overline{\mathscr{M}(f_k;s_k,q_k)})\\ &= (-1)^{d(s_k)-d(q_k)-1} o^*(\widetilde{\mathscr{D}}_{p_k}(f_k)) \wedge o^*(\widetilde{\mathscr{A}_{s_k}}(f_k)) \wedge o^*(\widetilde{\Sigma_k}) \wedge o^*(\overline{\mathscr{M}(f_k;s_k,q_k)})\\ &= (-1)^{d(s_k)-d(q_k)-1} o^*(\mathscr{M}_{\Gamma'}(\vec{f})) \wedge o^*(\overline{\mathscr{M}(f_k;s_k,q_k)}).\end{aligned}$$

This differs from the standard orientation by $(-1)^{d(s_k)-d(q_k)-1} = (-1)^{d(p_k)-d(s_k)}$.

(3) $X =$ 

. The induced orientation on the boundary is as in Lemma 3.17, which agrees with -1 of the standard orientation (3.9).

5. Proof of main theorem

5.1. Invariance on ordered 1-parameter family without i/i-intersections.

Lemma 5.1. *Suppose that* $\{f_s\}$ *is ordered and has no* i/i*-intersections over* $J = [s_0, s_1]$*. Then*

$$\widehat{Z}_{2k,3k}(\vec{f}_{s_0}) = \widehat{Z}_{2k,3k}(\vec{f}_{s_1}).$$

Proof. By Corollary 4.5 and the observation of §4.6, we have

$$\begin{aligned}&\partial\overline{\mathscr{M}}_\Gamma(\vec{f}_J)\\ &= -\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_1})\coprod\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0})\coprod\overline{\mathscr{M}}_{(d+d')\Gamma}(\vec{f}_J)\coprod(\partial^{\mathrm{hi}}\overline{\mathrm{Conf}}_n(M)\cap\overline{\mathscr{M}}_\Gamma(\vec{f}_J)),\end{aligned}$$

as oriented manifolds. Hence by Lemmas 3.20 and 3.21, we have

$$\begin{aligned}&\#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_1}) - \#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0})\\ &= \begin{cases}\#\overline{\mathscr{M}}_{(d+d')\Gamma}(\vec{f}_J) & \text{if } E(\Gamma)\neq\mathrm{Comp}(\Gamma)\\ \#\overline{\mathscr{M}}_{(d+d')\Gamma}(\vec{f}_J) + \#\overline{\mathscr{M}}_\Gamma^{\mathrm{local}}(-\mathrm{grad}_0 j^1\vec{f}_J) & \text{if } E(\Gamma)=\mathrm{Comp}(\Gamma)\end{cases}\end{aligned} \tag{5.1}$$

If Γ in the formula (5.1) is replaced with $\widetilde{\mathrm{Tr}}'_{\vec{g}}(\widetilde{\gamma}_{2k,3k})$, then the right hand side of (5.1) becomes

$$Z_{2k,3k}^{\mathrm{anomaly}}(j^3\vec{f}_J).$$

For the correction terms of $\widehat{Z}_{2k,3k}(\vec{f}_{s_0})$ and $\widehat{Z}_{2k,3k}(\vec{f}_{s_1})$, we may choose the same spin 4-manifold W. Then we choose generic sections $\vec{\rho}_W$ and $\vec{\rho'}_W$ of $(\mathscr{H}(T^vW))^m$ for the correction terms of $\widehat{Z}_{2k,3k}(\vec{f}_{s_0})$ and $\widehat{Z}_{2k,3k}(\vec{f}_{s_1})$ respectively, as in §3.10.3. Then by the proof of Proposition 3.29, we have

$$Z_{2k,3k}^{\mathrm{anomaly}}(j^3\vec{f}_J) - Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho}_W) + Z_{2k,3k}^{\mathrm{anomaly}}(\vec{\rho'}_W) = 0.$$

This completes the proof. □

5.2. **Invariance at level exchange point.**

Lemma 5.2. *Suppose that* $s_0\in J$ *is a level exchange point for the 1-parameter family* $\{f_s\}_{s\in J}$. *Then*

$$\widehat{Z}_{2k,3k}(\vec{f}_{s_0-\varepsilon}) = \widehat{Z}_{2k,3k}(\vec{f}_{s_0+\varepsilon}).$$

The proof is the same as Lemma 5.1.

5.3. **Invariance at birth/death point.** We say that a birth/death point v in a 1-parameter family $\{(f_s,\mu_s)\}_{s\in[0,1]}$, say at $s=s_0$, is *almost independent* if on a neighborhood of s_0 in $[0,1]$ the descending and ascending manifolds of v are disjoint from all the other descending and ascending manifolds of critical points except those of the pair $(\widetilde{a}(s),\widetilde{b}(s))$ chosen in Lemma 4.2.

Lemma 5.3 ([HW], page 62). *A 1-parameter family* $\{(f_s,\mu_s)\}_{s\in[0,1]}$ *of pairs of generalized Morse functions and metrics on* M *can be deformed so that every birth/death points are almost independent.*

Lemma 5.4. *Suppose that* $s_0\in J$ *is a parameter on which an almost independent birth/death point* v *occurs. Then*

$$\widehat{Z}_{2k,3k}(\vec{f}_{s_0-\varepsilon}) = \widehat{Z}_{2k,3k}(\vec{f}_{s_0+\varepsilon}).$$

Proof. We prove the lemma only for death point since the case of birth point is symmetric. If (p,q) is the critical point pair at $s=s_0-\varepsilon$ that disappears on $s\geq s_0$, then the Morse complex at $s=s_0-\varepsilon$ is the direct sum $\widetilde{C}_*^{(s_0+\varepsilon)}\oplus C_*^{\mathrm{elem}}$, where $\widetilde{C}_*^{(s_0+\varepsilon)} = (\widetilde{C}_*^{(s_0+\varepsilon)},\widetilde{\partial}^{(s_0+\varepsilon)})$ is the reduced Morse complex at $s=s_0+\varepsilon$ and

$$C_*^{\mathrm{elem}} = \{0\to C_{i+1}^{\mathrm{elem}}=\mathbb{Z}^{\{p\}}\overset{\approx}{\to} C_i^{\mathrm{elem}}=\mathbb{Z}^{\{q\}}\to 0\}.$$

Choose a combinatorial propagator g of $\widetilde{C}_*^{(s_0+\varepsilon)}$. The acyclic complex C_*^{elem} has a unique combinatorial propagator g^{elem}, defined by $g^{\mathrm{elem}}(q) = p$. We consider g and g^{elem} as homogeneous degree 1 maps of $\widetilde{C}_*^{(s_0+\varepsilon)} \oplus C_*^{\mathrm{elem}}$ by setting $g(C_*^{\mathrm{elem}}) = 0$ and $g^{\mathrm{elem}}(\widetilde{C}_*^{(s_0+\varepsilon)}) = 0$. Then one can check that $g' = g + g^{\mathrm{elem}}$ is a combinatorial propagator for $\widetilde{C}_*^{(s_0+\varepsilon)} \oplus C_*^{\mathrm{elem}}$.

We need only to check the identity of the lemma in the case where a gluing of trajectories happens at v. Suppose that the separated edge labeled by k in a graph flow of $\Gamma(p,q)_k \in \mathscr{G}_{2k,3k}(\vec{C}^{(s_0-\varepsilon)})$ converges to a broken edge as $s \to s_0$ and that p and q converges to v. Then by Corollary 4.9 we have

$$\begin{aligned}
&\widetilde{\mathrm{Tr}}_{\cdots,g',\cdots}\big[[\Gamma(p,q)_k]\big] \cdot \#\overline{\mathscr{M}}_{\Gamma(p,q)_k}(\vec{f}_{s_0-\varepsilon}) + \widetilde{\mathrm{Tr}}_{\cdots,g',\cdots}\big[[\Gamma(\emptyset,\emptyset)_k]\big] \cdot \#\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_k}(\vec{f}_{s_0-\varepsilon})\\
&= \widetilde{\mathrm{Tr}}_{\cdots,g,\cdots}\big[[\Gamma(\emptyset,\emptyset)_k]\big]\Big(\#\overline{\mathscr{M}}_{\Gamma(p,q)_k}(\vec{f}_{s_0-\varepsilon}) + \#\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_k}(\vec{f}_{s_0-\varepsilon})\Big)\\
&= \widetilde{\mathrm{Tr}}_{\cdots,g,\cdots}\big[[\Gamma(\emptyset,\emptyset)_k]\big] \cdot \#\overline{\mathscr{M}}_{\Gamma(\emptyset,\emptyset)_k}(\vec{f}_{s_0+\varepsilon}).
\end{aligned}$$

The proof of the invariance of the other terms in $\widehat{Z}_{2k,3k}(\vec{f}_{s_0-\varepsilon})$ is the same as Lemma 5.1 and 5.2 since v is almost independent. □

5.4. **Invariance at i/i-intersection point.** Suppose that a gradient i/i-intersection between critical points (loci) p and q occurs at $s = s_0$. For a small number $\varepsilon > 0$, the underlying $\mathbb{Z}$-modules of $\widetilde{C}_*^{(s_0-\varepsilon)}$ and $\widetilde{C}_*^{(s_0+\varepsilon)}$ are the same. We put $J = [s_0-\varepsilon, s_0+\varepsilon]$, $\widetilde{C}_*^{(J)} = \widetilde{C}_*^{(s_0-\varepsilon)} = \widetilde{C}_*^{(s_0+\varepsilon)}$ and $P_*^{(J)} = P_*^{(s_0-\varepsilon)} = P_*^{(s_0+\varepsilon)}$. Let $h : \widetilde{C}_*^{(J)} \to \widetilde{C}_*^{(J)}$ be the homomorphism of homogeneous degree 0, defined for each critical point (locus) $a \in P_i^{(J)}$ by

$$\begin{aligned}
h(a) &= \sum_{b \in P_i^{(J)}} \#\overline{\mathscr{M}(f_J;a,b)} \cdot b,\\
\overline{\mathscr{M}(f_J;a,b)} &= \mathscr{M}(f_J;a,b) \pitchfork \widetilde{L}_a,\\
\mathscr{M}(f_J;a,b) &= \widetilde{\mathscr{D}_a}(f_J) \pitchfork \widetilde{\mathscr{A}_b}(f_J).
\end{aligned}$$

We denote the boundary operators of $\widetilde{C}_*^{(s_0-\varepsilon)}$ and $\widetilde{C}_*^{(s_0+\varepsilon)}$ by ∂ and ∂' respectively.

Lemma 5.5. *Under the assumption above, we have*

$$\partial' - \partial = \partial' h - h\partial = \partial h - h\partial',$$

or equivalently, $\partial'(1-h) = (1-h)\partial$ *and* $\partial(1+h) = (1+h)\partial'$.

More explicitly, if $p, q \in P_i^{(J)}$ and if $r \in P_{i+1}^{(J)}$ is any other critical point of index $i+1$, then it follows from Lemma 5.5 that

$$h(x) = \begin{cases} \pm q & \text{if } x = p\\ 0 & \text{otherwise}\end{cases}$$

and that the difference of ∂' and ∂ is given by

$$\begin{cases} \partial' p = \partial p \pm \partial q\\ \partial' r = \partial r \mp \partial_{rp} \cdot q = \partial r \mp \partial'_{rp} \cdot q\\ \partial' q = \partial q \end{cases} \tag{5.2}$$

In particular, we have $h^2 = 0$.

Proof of Lemma 5.5. We know from Corollary 4.5 what happens at s_0, so it suffices to check that the signs are correct. We may assume without loss of generality that only the ascending manifold of q changes as s moves within J if ε is small enough. So we may replace the problem with the intersection orientations in a single fiber in which the ascending manifold of q moves around, depending on a parameter $s \in J$.

There is a small neighborhood of $\mathscr{F}_p\overline{\mathscr{M}}_2(f_J;r,q) \cong \overline{\mathscr{M}(f_{s_0};r,p)} \times \overline{\mathscr{M}(f_J;p,q)}$ that is a subspace of $U_a \times \overline{L}_p$ for some small open ball U_a in a Morse model M_p around p (for the quadratic form $-x_1^2-\cdots-x_i^2+x_{i+1}^2+\cdots+x_d^2$). By an analogue of Lemma 3.16, the normal orientation $o^*(\overline{\mathscr{M}}_2(f_J;r,q))_{(x,x')}$ at $(x,x') \in U_a \times \overline{L}_p$ induces an orientation

$$\begin{aligned}(5.3)\qquad & o^*(\mathscr{D}_r(f_{s_0}))_x \wedge (-1)^{d(p)-(d(q)+1)} o^*(\mathscr{A}_p(f_{s_0}))_x \wedge o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'} \\ & = -o^*(\mathscr{M}(f_{s_0};r,p))_x \wedge o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'}\end{aligned}$$

on the space of (partially parametrized) broken trajectories. We assume without loss of generality that $U_a \cap \overline{L'_p} \neq \emptyset$ and that the points x and $(0,\ldots,0,\sqrt{\varepsilon}) \in M_p$ lie on the same gradient trajectory. Since $o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'}$ is a d-form, $o^*(\overline{L'_p})_x = dx_d$ and $o^*(\overline{L'_p} \times \mathbb{R}^d)_{(x,x')} = *\iota\big(\frac{\partial}{\partial x_d}\big)dx_1 \wedge \cdots \wedge dx_d \wedge dx'_1 \wedge \cdots \wedge dx'_d = (-1)^d dx_d$, the orientation (5.3) induces the following orientation on the intersection with $\overline{L'_p} \times \overline{L}_p$:

$$\begin{aligned}(5.4)\qquad & -o^*(\mathscr{M}(f_{s_0};r,p))_x \wedge o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'} \wedge (-1)^d dx_d \\ & = -o^*(\mathscr{M}(f_{s_0};r,p))_x \wedge dx_d \wedge o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'} \\ & = -o^*(\overline{\mathscr{M}(f_{s_0};r,p)})_x \wedge o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'} \\ & = -o^*(\overline{\mathscr{M}(f_{s_0-\varepsilon};r,p)})_x \wedge o^*(\overline{\mathscr{M}(f_J;p,q)})_{x'}\end{aligned}$$

This gives the sign of the incidence coefficient of q in $h\partial(r)$.

Similarly, for a critical point r' of index $i-1$, a small neighborhood of $\mathscr{F}_q\overline{\mathscr{M}}_2(f_J;p,r') \cong \overline{\mathscr{M}(f_J;p,q)} \times \overline{\mathscr{M}(f_{s_0};q,r')}$ can be considered as a subspace of $\overline{L'_q} \times U_a$ for some small open ball U_a in a Morse model M_q around q (for the quadratic form $-x_1^2-\cdots-x_i^2+x_{i+1}^2+\cdots+x_d^2$). Then by an analogue of Lemma 3.15, the normal orientation $o^*(\overline{\mathscr{M}}_2(f_J;p,r'))_{(x,x')}$ at $(x,x') \in \overline{L'_q} \times U_a$ induces an orientation

$$\begin{aligned}(5.5)\qquad & o^*(\overline{\mathscr{M}(f_J;p,q)})_x \wedge o^*(\mathscr{D}_q(f_{s_0}))_{x'} \wedge o^*(\mathscr{A}_{r'}(f_{s_0}))_{x'} \\ & = o^*(\overline{\mathscr{M}(f_J;p,q)})_x \wedge o^*(\mathscr{M}(f_{s_0};q,r'))_{x'}\end{aligned}$$

on the space of (partially parametrized) broken trajectories. We assume without loss of generality that $U_a \cap \overline{L}_q \neq \emptyset$ and that the points x' and $(\sqrt{\varepsilon},0,\ldots,0) \in M_q$ lie on the same gradient trajectory. Since $o^*(\overline{L}_q)_{x'} = o^*(\mathbb{R}^d \times \overline{L}_q)_{(x,x')} = -dx'_1$, (5.5) induces the following orientation on the intersection with $\overline{L'_q} \times \overline{L}_q$:

$$\begin{aligned}(5.6)\qquad & o^*(\overline{\mathscr{M}(f_J;p,q)})_x \wedge o^*(\mathscr{M}(f_{s_0};q,r'))_{x'} \wedge (-dx'_1) \\ & = o^*(\overline{\mathscr{M}(f_J;p,q)})_x \wedge o^*(\overline{\mathscr{M}(f_{s_0};q,r')})_{x'} \\ & = o^*(\overline{\mathscr{M}(f_J;p,q)})_x \wedge o^*(\overline{\mathscr{M}(f_{s_0+\varepsilon};q,r')})_{x'}\end{aligned}$$

This gives the sign of the incidence coefficient of r' in $\partial' h(p)$.

The induced orientations (5.4) and (5.6) give the desired identities $\partial - \partial' - h\partial + \partial' h = 0$ and $\partial - \partial' - h\partial' + \partial h = 0$. $\square$

Choose a combinatorial propagator g for $(\widetilde{C}_*^{(s_0-\varepsilon)},\partial)$. Then one can check that the endomorphism

$$g'=(1-h)\circ g\circ(1+h)\in \mathrm{End}_1(\widetilde{C}_*^{(s_0+\varepsilon)})$$

is a combinatorial propagator for $(\widetilde{C}_*^{(s_0+\varepsilon)},\partial')$. One can check that $hgh=0$ and hence

$$g'-g=gh-hg.$$

We put $\vec{C}=(\widetilde{C}_*^{(s_0-\varepsilon)},\widetilde{C}_*^{(2)},\ldots,\widetilde{C}_*^{(m)})$ and $\vec{C}'=(\widetilde{C}_*^{(s_0+\varepsilon)},\widetilde{C}_*^{(2)},\ldots,\widetilde{C}_*^{(m)})$. We define the operator $d'':\mathscr{G}_{n,m,\vec{\eta}}(\vec{C})\to\mathscr{G}_{n,m-1,d''\vec{\eta}}(\vec{C})$ by $\sum_{e\in\mathrm{Se}(\Gamma)}d''_e$, where

d''_e 

$= \displaystyle\sum_{\substack{r_i\in P_*^{(i)}\\ d(r_i)=d(p_i)}}$ 

$+ \displaystyle\sum_{\substack{s_i\in P_*^{(i)}\\ d(s_i)=d(q_i)}} (-1)^{d(p_i)-d(s_i)}$ 

$(\beta(i)=e)$

and $\mathrm{or}(d''_e\Gamma)$ is the induced one. The reason for the sign has been explained in §4.6.

Lemma 5.6. *Suppose that* $s_0\in J$ *is a point on which a gradient i/i-intersection between critical points (loci)* p *and* q *occurs in a 1-parameter family* $\{(f_s,\mu_s)\}_{s\in J}$. *Then*

$$\widehat{Z}_{2k,3k}(\vec{f}_{s_0-\varepsilon})=\widehat{Z}_{2k,3k}(\vec{f}_{s_0+\varepsilon}).$$

Proof. As oriented manifolds, we may assume that

$$\partial\overline{\mathscr{M}}_\Gamma(\vec{f}_J)=-\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0+\varepsilon})\coprod\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon})\coprod\overline{\mathscr{M}}_{d''\Gamma}(\vec{f}_J)$$

if ε is small enough. Hence we have

$$\#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0+\varepsilon})-\#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon})-\#\overline{\mathscr{M}}_{d''\Gamma}(\vec{f}_J)=0.$$

We put $\vec{g} = (g, g_2, \dots, g_m)$ and $\vec{g}' = (g', g_2, \dots, g_m)$. We have

$$\begin{aligned}
&\sum_{\Gamma \in \mathscr{G}_{2k,3k}(\vec{C})} \widetilde{\mathrm{Tr}}_{\vec{g}'}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0+\varepsilon}) - \sum_{\Gamma \in \mathscr{G}_{2k,3k}(\vec{C}')} \widetilde{\mathrm{Tr}}_{\vec{g}}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon}) \\
&= \sum_\Gamma \widetilde{\mathrm{Tr}}_{\vec{g}'}\big[[\Gamma]\big] \cdot \Big(\#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon}) + \#\overline{\mathscr{M}}_{d''\Gamma}(\vec{f}_J)\Big) - \sum_\Gamma \widetilde{\mathrm{Tr}}_{\vec{g}}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon}) \\
&= \sum_\Gamma \widetilde{\mathrm{Tr}}_{g'-g,\dots}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon}) + \sum_\Gamma \widetilde{\mathrm{Tr}}_{\vec{g}'}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_{d''\Gamma}(\vec{f}_J) \\
&= \sum_\Gamma \widetilde{\mathrm{Tr}}_{gh-hg,\dots}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_\Gamma(\vec{f}_{s_0-\varepsilon}) + \sum_\Gamma \widetilde{\mathrm{Tr}}_{\vec{g}'}\big[[\Gamma]\big] \cdot \#\overline{\mathscr{M}}_{d''\Gamma}(\vec{f}_J) \\
&= \widetilde{\mathrm{Tr}}_{g,\dots}\Big[\sum_{\substack{p_1,q_1 \\ d(p_1)=d(q_1)+1}} \sum_{\Gamma(p_1,q_1)_1} [\Gamma(p_1,q_1)_1] \cdot \#\overline{\mathscr{M}}_{(\Gamma * h - h * \Gamma)(p_1,q_1)_1}(\vec{f}_J)\Big] \\
&\qquad + \widetilde{\mathrm{Tr}}_{\vec{g}'}\Big[\sum_{\substack{p_1,q_1 \\ d(p_1)=d(q_1)+1}} \sum_{\Gamma(p_1,q_1)_1} [\Gamma(p_1,q_1)_1] \cdot \#\overline{\mathscr{M}}_{d''\Gamma(p_1,q_1)_1}(\vec{f}_J)\Big] \\
&= \sum_{p_1,q_1} \sum_{\Gamma(p_1,q_1)_1} \widetilde{\mathrm{Tr}}_{g'-g,\dots}\big[[\Gamma(p_1,q_1)_1]\big] \cdot \#\overline{\mathscr{M}}_{d''\Gamma(p_1,q_1)_1}(\vec{f}_J) \\
&= \sum_{p_1,q_1} \sum_{\Gamma(p_1,q_1)_1} \widetilde{\mathrm{Tr}}_{gh-hg,\dots}\big[[\Gamma(p_1,q_1)_1]\big] \cdot \#\overline{\mathscr{M}}_{d''\Gamma(p_1,q_1)_1}(\vec{f}_J) \\
&= -\sum_{p_1,q_1} \sum_{\Gamma(p_1,q_1)_1} \widetilde{\mathrm{Tr}}_{g,\dots}\big[[\Gamma(p_1,q_1)_1]\big] \cdot \#\overline{\mathscr{M}}_{d''d''\Gamma(p_1,q_1)_1}(\vec{f}_J) \\
&= 0
\end{aligned}$$

since $h^2 = 0$ and $hgh = 0$. This completes the proof. □

Proof of Theorem 3.32. Lemmas 5.1, 5.2, 5.4, 5.6 show that $\widehat{Z}_{2k,3k}(\vec{f}_s)$ is invariant under all possible bifurcations listed in §4.1. This completes the proof. □

APPENDIX A. ORIENTATIONS ON MANIFOLDS AND THEIR INTERSECTIONS

A.1. **Orientation on a manifold.** For a d-dimensional orientable manifold M, we will represent an orientation on M by a nowhere vanishing d-form of $\Omega^d_{\mathrm{dR}}(M)$ and denote by $o(M)$. If M is a submanifold of an oriented d'-dimensional manifold N, then we define $o(M)$ from an orientation $o^*(M)$ of the normal bundle of M by the rule

$$o(M) \wedge o^*(M) \sim o(N).$$

Note that $o^*(M)$ is defined canonically by the Hodge star operator: $o^*(M) = *o(M)$.

A.2. **Induced orientation.** If M has boundary ∂M, we provide an induced orientation on ∂M from $o(M)$ as follows: let n be an inward normal vector field on ∂M, then we define

$$o(\partial M)_x = \iota(n_x) o(M)_x.$$

In other words, if n_x^* is the dual of n_x with respect to the metric and if $o(M)_x = \varepsilon\, n_x^* \wedge \alpha_x$ for $\alpha_x \in \Omega^{d-1}_{\mathrm{dR}}(M)$ and $\varepsilon \in \{1, -1\}$, then $o(\partial M)_x = \varepsilon \alpha_x$. This change induces a change on normal orientation $o^*(M)$ as follows: $o^*(M)_x \mapsto (-1)^{d'-1} o^*(M)_x \wedge$

n_x^*. So we have

$$o^*(\partial M)_x = (-1)^{d'-1} o^*(M)_x \wedge n_x^*. \tag{A.1}$$

Example A.1. Suppose $M = \{(x_1, \ldots, x_i, 0, \ldots, 0) \in \mathbb{R}^d;\, x_1 \geq 0\}$ and $N = \mathbb{R}^d$. We assume that $o(M)_x = dx_1 \wedge \cdots \wedge dx_i$ and $o(\mathbb{R}^d)_x = dx_1 \wedge \cdots \wedge dx_d$, then $o(\partial M)_x = \iota\big(\frac{\partial}{\partial x_1}\big) o(M)_x = dx_2 \wedge \cdots \wedge dx_i$. On the other hand, we have $o^*(M)_x = dx_{i+1} \wedge \cdots \wedge dx_d$ and $o^*(\partial M)_x = (-1)^{i-1} dx_1 \wedge dx_{i+1} \wedge \cdots \wedge dx_d = (-1)^{d-1} dx_{i+1} \wedge \cdots \wedge dx_d \wedge dx_1$.

We equip each level surface L of a Morse function $f : M \to \mathbb{R}$ by the rule

$$o(L)_x = \iota(\operatorname{grad} f)_x o(M)_x.$$

A.3. **Orientation of intersection.** Suppose M and M' are two oriented submanifolds of N of dimension i and i' that intersect transversally. The transversality implies that at an intersection point x, the form $o^*(M)_x \wedge o^*(M')_x$ is a non-trivial $(2d - i - i')$-form. We define

$$o^*(M \pitchfork M')_x = o^*(M)_x \wedge o^*(M')_x.$$

Acknowledgments. I would like to thank Professor Kenji Fukaya for encouraging me to write my result for publication. I would also like to thank Professor Masamichi Takase for valuable comments on invariants of spin 4-manifolds. I am supported by JSPS Grant-in-Aid for Young Scientists (B) 70467447.

References

[AS] S. Axelrod, I. M. Singer, *Chern–Simons perturbation theory*, in Proceedings of the XXth DGM Conference, Catto S., Rocha A. (eds.), pp. 3–45, World Scientific, Singapore, 1992.

[BC] M. Betz, R. Cohen, *Graph Moduli Spaces and Cohomology Operations*, Turkish J. Math. **18** (1995), 23–41.

[Ce] J. Cerf, *La stratification naturelle des espaces de fonctions différentiables réelles et le théorème de la pseudo-isotopie*, Publ. Math. I.H.É.S. **39** (1970), 5–173.

[Fuk93] K. Fukaya, *Morse homotopy, A^∞-category and Floer homologies*, in Proceedings of GARC Workshop on Geometry and Topology, ed. H. J. Kim, Seoul National University, Korea, 1993.

[Fuk96] K. Fukaya, *Morse Homotopy and Chern–Simons Perturbation Theory*, Comm. Math. Phys. **181** (1996), 37–90.

[Fut] M. Futaki, Master thesis, Univ. of Tokyo.

[FO] K. Fukaya, Y-G. O, *Zero-loop open strings in the cotangent bundle and Morse homotopy*, Asian J. Math. **1** (1997), no. 1, 96–180.

[FM] W. Fulton, R. MacPherson, *A compactification of configuration spaces*, Ann. of Math. **139** (1994), 183–225.

[HW] A. Hatcher, J. Wagoner, *Pseudo-isotopies of compact manifolds*, Astérisque, No. 6. Société Mathématique de France, Paris, 1973. i+275 pp.

[Ig1] K. Igusa, *The space of framed functions*, Trans. Amer. Math. Soc. **301** (1987), no. 2, 431–477.

[Ig2] K. Igusa, *The stability theorem for smooth pseudoisotopies*, K-Theory **2** (1988), 1–355.

[KM] R. Kirby, P. Melvin, *Canonical framings for 3-manifolds*, Proceedings of 6th Gökova Geometry-Topology Conference, Turkish J. Math. **23** (1999), no. 1, 89–115.

[Ko] M. Kontsevich, *Feynman diagrams and low-dimensional topology*, First European Cogress of Mathematics, Vol. II (Paris, 1992), Progr. Math. **120** (Birkhauser, Basel, 1994), 97–121.

[KT] G. Kuperberg, D. Thurston, *Perturbative 3-manifold invariants by cut-and-paste topology*, preprint, arXiv:math/9912167.

[Les] C. Lescop, *Splitting formulae for the Kontsevich–Kuperberg–Thurston invariant of rational homology 3-spheres*, math.GT/0411431.

[Oh96] T. Ohtsuki, *Finite type invariants of integral homology 3-spheres*, J. Knot Theory Ramifications **5** (1996), 101–115.

[Wa] T. Watanabe, *Obstructions for* π_1 *of the space of pseudo-isotopies of disks in non-stable range*, preprint.

[Wa2] T. Watanabe, *Higher order generalization of Fukaya's Morse homotopy invariant of 3-manifolds II*, in preparation.

[Wi] E. Witten, *Chern–Simons gauge theory as a string theory*, The Floer memorial volume, 637–678, Progr. Math., 133, Birkhäuser, Basel, 1995.

Department of Mathematics, Hokkaido University, Kita 10, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810, Japan

E-mail address: `tadayuki@math.sci.hokudai.ac.jp`